

Apr 18th, 10:50 AM - 11:10 AM

Acknowledging Sampling Bias in Species Distribution Modelling: Predicting *Botrychium* sp. Habitat in Lincoln County, Montana

Annalisa Ingegno
ai140843@umconnect.umt.edu

Follow this and additional works at: <http://scholarworks.umt.edu/gsrc>

Annalisa Ingegno, "Acknowledging Sampling Bias in Species Distribution Modelling: Predicting *Botrychium* sp. Habitat in Lincoln County, Montana" (April 18, 2015). *UM Graduate Conference (GradCon)*. Paper 3.
<http://scholarworks.umt.edu/gsrc/2015/oralpres1d/3>

This Oral Presentation is brought to you for free and open access by ScholarWorks at University of Montana. It has been accepted for inclusion in UM Graduate Conference (GradCon) by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact scholarworks@mail.lib.umt.edu.

INTRODUCTION : Over 70 % of *Botrychium* in Montana are considered “Species of Concern” by the Montana Natural Heritage Program, meaning they are in jeopardy of extirpation due to habitat destruction, limited range, or small population size.¹² Thus, developing an effective conservation plan for Montana that maintains current *Botrychium* sp. diversity and preserves their distribution is critical.^{5,4} Unfortunately, *Botrychium* sp. periodically abstain from producing aboveground fronds due to their symbiosis with arbuscular mycorrhiza, complicating long-term data collection strategies.^{9,8,4} Ergo, *Botrychium* sp. make an ideal candidate for a species distribution model (SDM) because it can provide statistical evidence linking observed distribution with environmental variables.^{7,3,2} Frequently applied in both conservation and management activities, an SDM can predict potential distribution over an area which is pivotal to conservation planning.^{5,13} Studies have yet to employ this technique for *Botrychium* sp. and thus, there is no way to assess status or identify threats, putting *Botrychium* sp. at serious risk.^{5,12} *An SDM was used to identify environmental predictors for Botrychium sp. habitat in order to construct a predictive distribution map for Botrychium sp. in Lincoln County, Montana.*

BACKGROUND : In fall 2014, a preliminary feasibility analysis focused on evaluating correction techniques to mitigate the sampling bias found in *Botrychium* sp. occurrence data from Lincoln County, Montana using MaxEnt. Maximum Entropy (MaxEnt) is an SDM that associates the spatial location of occurrence data with environmental variables to predict habitat suitability at unsurveyed locations.¹³ Sampling bias arises when observation data is preferentially collected from certain areas because they are for example, more convenient to reach.^{10,2,6} Eight explanatory variables were considered as input to the model: elevation, aspect, slope, soils, geology, mean May precipitation, mean June temperature, and land cover.^{4,14} Using techniques assembled from similar studies, sampling bias correction schemes were created that split the data into two categories of locational uncertainty, subsampled the data at two resolutions, or restricted the background sampling extent to the known distribution of *Botrychium* sp.^{3,10,2,6} Sampling bias was said to be resolved if the dataset exhibited a random distribution after application of a correction technique. Three optimal datasets were produced from the corrective procedures: points split based on locational uncertainty and then subsampled, points split based on locational uncertainty alone, and the complete *Botrychium* sp. dataset. The split and subsampled dataset performed poorly, creating a suitability map that grossly over-predicted potential habitat. Because the original sampling bias was so strong, this correction produced a subset of data that was no longer representative of the entire study area. Consequently, future analyses should continue to experiment with subsampling but refrain from splitting data into locational uncertainty classes.

RESEARCH PLAN :

(1) *Correction of sampling bias in observation data:* (i) Splitting - data will not be split based upon locational uncertainty. Further scrutiny will be given to the process of eliminating redundant points; relevant data was incidentally removed during this process previously. (ii) Subsampling - there exist two main clusters of points in the survey area; limiting subsampling to only the clustered regions will be further examined. (iii) Background Extent - quantification of excluding unsuitable habitat like lakes, rivers, and roads, and high elevation sites will be examined.

(2) *Specification of environmental variables:* Categories of environmental variables were assembled ranging from standard climatic variables, edaphic variables, and climatic variables that influence edaphic variables.

(3) *Testing of MaxEnt's internal settings*: MaxEnt has the ability to modify the number of runs, the type of evaluation procedure, the number of background points sampled, and more. These can considerably alter results.¹³

CONCLUSION : This analysis is innovative in its application of an SDM to address deficiencies in knowledge about an enigmatic subgenus of Ophioglossaceae known as *Botrychium*. It will be one of the first studies to provide statistical support for linking environmental variables to observed *Botrychium* sp. habitat. Furthermore, this research will enhance our understanding of the ecological mechanisms behind symbiotic relationships between plants and arbuscular mycorrhiza. It will fill an important knowledge gap on the current distribution of *Botrychium* sp. in Lincoln County, MT and has potential to be expanded to other counties in Montana as well. It will help us monitor the health of this subgenus through changing disturbance and climatic regimes and create a conservation strategy that is relevant for maintaining current populations across the state.

LITERATURE CITED

- ¹ Ahlenslager, Kathy, and Laura Potash. "Conservation Assessment for 13 Species of Moonworts (*Botrychium* Swartz Subgenus *Botrychium*)." Report Submitted to USDA Forest Service Region 6 and USDI Bureau of Land Management, Oregon and Washington. April 18, 2007. ² Boria, Robert A., Link E. Olson, Steven M. Goodman, and Robert P. Anderson. "Spatial Filtering to Reduce Sampling Bias Can Improve the Performance of Ecological Niche Models." *Ecological Modelling*, 275(2014): 73-77.
- ³ Burbach, Thor. "The Influence of Environmental Variables on Predicting Rare-Plant Habitat in the Nez Perce National Forest." Master's Thesis, University of Montana, 2011. ⁴ Farrar, Donald R. "Systematics of moonworts *Botrychium* subgenus *Botrychium*." Iowa : Iowa State University, Department of Ecology, Evolution, and Organismal Biology, 2006. ⁵ Fielding, Alan H. and John F. Bell. "A Review of Methods for the Assessment of Prediction Errors in Conservation Presence/Absence Models." *Environmental Conservation*, 24(1997): 38-49. ⁶ Fourcade, Yoan, Jan O. Engler, Dennis Rodder, and Jean Secondi. "Mapping Species Distributions with MAXENT Using a Geographically Based Sample of Presence Data : A Performance Assessment of Methods for Correcting Sampling Bias." *PLOS ONE* 9, 3(2014) : e97122. ⁷ Guisan, Antoine, and Niklaus E. Zimmermann. "Predictive habitat distribution models in ecology." *Ecological Modelling*, 135(2000): 147-186.
- ⁸ Johnson-Groh, Cindy L., and Jennifer M. Lee. "Phenology and Demography of Two Species of *Botrychium* (Ophioglossaceae)." *American Journal of Botany*, 89(2002a): 1624-1633. ⁹ Johnson-Groh, Cindy, Chandra Riedel, Laura Schoessler, and Krissa Skogen. "Belowground Distribution and Abundance of *Botrychium* Gametophytes and Juvenile Sporophytes." *American Fern Journal*, 92(2002b): 80-92. ¹⁰ Kramer-Schadt, Stephanie, Jürgen Niedballa, John D. Pilgrim, Boris Schröder, Jana Lindenborn, Vanessa Reinfelder, Milena Stillfried, Ilja Heckmann, Anne K. Scharf, Dave M. Augeri, Susan M. Cheyne, Andrew J. Hearn, Joanna Ross, David W. Macdonald, John Mathai, James Eaton, Andrew J. Marshall, Gono Semiadi, Rustam, Henry Bernard, Raymond Alfred, Hiromitsu Samejima, J. W. Duckworth, Christine Breitenmoser-Wuersten, Jerrold L. Belant, Heribert Hofer, and Andreas Wilting. "The Importance of Correcting for Sampling Bias in MaxEnt Species Distribution Models." *Diversity and Distributions*, 19 no. 11(2013): 1366–1379. ¹¹ Lesica, Peter, and Kathleen Ahlenslager. "Demography and Life History of Three Sympatric Species of *Botrychium* subg. *Botrychium* in Waterton Lakes National Park, Alberta." *Canadian Journal of Botany*, 74(1996): 538-543. ¹² Montana Natural Heritage Program. "Montana Plant Species of Concern Report." State of Montana, 2014. Available at : <http://mtnhp.org/SpeciesOfConcern/?AorP=p>. ¹³ Phillips, Steven J., Robert P. Anderson, and Robert E. Schapire. "Maximum entropy modeling of species geographic distributions." *Ecological Modelling*, 190(2006): 231-259.
- ¹⁴ Roe-Anderson, Susan M., and Darlene Southworth. "Microsite Factors and Spore Dispersal Limit Obligate Mycorrhizal Fern Distribution: Habitat Islands of *Botrychium* pumicola (Ophioglossaceae)." *American Fern Journal*, 103(2013): 1-20.