

University of Montana

## ScholarWorks at University of Montana

---

Health and Wellness

Rural Institute for Inclusive Communities

---

8-2014

### Pain and Participation

Catherine Ipsen

*University of Montana Rural Institute - Research and Training Center on Disability in Rural Communities*

Tannis Hargrove

*University of Montana Rural Institute - Research and Training Center on Disability in Rural Communities*

Naomi Kimbell

*University of Montana Rural Institute - Research and Training Center on Disability in Rural Communities*

University of Montana Rural Institute

[scholarworks-reports@mso.umt.edu](mailto:scholarworks-reports@mso.umt.edu)

Follow this and additional works at: [https://scholarworks.umt.edu/ruralinst\\_health\\_wellness](https://scholarworks.umt.edu/ruralinst_health_wellness)



Part of the [Community Health and Preventive Medicine Commons](#)

**Let us know how access to this document benefits you.**

---

#### Recommended Citation

Ipsen, C., Hargrove, T., & Kimbell, N. (2014). Pain and Participation. Missoula: The University of Montana Rural Institute.

This Research Summary is brought to you for free and open access by the Rural Institute for Inclusive Communities at ScholarWorks at University of Montana. It has been accepted for inclusion in Health and Wellness by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact [scholarworks@mso.umt.edu](mailto:scholarworks@mso.umt.edu).

# Research Summary

## Pain and Participation

The objective of the Pain Interference Patterns research project is to understand how pain and environmental conditions, like access barriers, affect community participation of people who report physical, sensory and cognitive impairment. The information will be used to develop strategies for increasing participation in rural communities.

People with disabilities often face barriers to full participation in home, school, work and community activities. Barriers to participation include a variety of factors including but not limited to:

- Inaccessible transportation
- Inaccessible buildings and public spaces
- Health conditions like depression, circulation problems and pain

Because the majority of individuals with a disability report experiencing pain as a secondary condition, we wanted to know how pain impacted participation in life activities. To help answer this question, we asked individuals with disabilities to complete four surveys over 18 months about their pain levels, environmental barriers, and participation in daily activities.

### Research Participants

Of the 525 people who participated in this research study, most reported some kind of impairment, from hearing and vision loss to significant mobility loss. As a whole, the group reported an average daily pain level of 4.68 on a 0-10 pain scale (see Figure 1).




Figure 1. Pain Rating Scale

RTC:Rural  
52 Corbin Hall  
University of Montana  
Missoula, MT 59812  
Toll Free: 888.268.2743  
Fax: 406.243.2349  
TTY: 406.243.4200

rtcrural@mso.umt.edu  
rtc.ruralinstitute.umt.edu  
Alternative formats available

*RTC:Rural conducts research, education, and training to improve the ability of people with disabilities to live full and independent lives in rural communities.*

# Preliminary Results

Based on responses from the first survey, we learned that:

- Community participation can be broken into three categories:
  - Medical, including doctor, other healthcare and pharmacy visits
  - Shopping, including grocery stores, shopping malls, and box stores
  - Free time, including religious activities, socializing outside the home, community activities, exercise, and entertainment
- Individuals who participated more frequently in community activities reported higher rates of pleasure and happiness.
- Married people were less likely to participate in free time activities than their single counterparts.
- People who managed pain with rest had less free time activities.
- People with high pain levels had less medical and free time activities.
- The way people thought about pain was more of a barrier than pain itself—this is called catastrophic thinking about pain.

## Conclusion

Although we are still analyzing data, we are gaining a better understanding of how pain interferes with participation. For more technical information about the participation categories and analyses, please see our research report *Pain Interference Patterns* at [rtc.ruralinstitute.umt.edu/rtcBlog/wp-content/uploads/PIP.pdf](http://rtc.ruralinstitute.umt.edu/rtcBlog/wp-content/uploads/PIP.pdf) and look for more research reports in the future.

**Prepared by:** Catherine Ipsen, Tannis Hargrove and Naomi Kimbell

**Primary Researchers:** Craig Ravesloot, Catherine Ipsen, Tannis Hargrove, Lillie Greiman and Sandra Skogley

**Suggested citation:** Ipsen, C., Hargrove, T., & Kimbell, N. (2014). *Pain and Participation*. Missoula: The University of Montana Rural Institute.

**For additional information please contact:**

Research and Training Center on Disability in Rural Communities; University of Montana Rural Institute; 52 Corbin Hall, Missoula, MT 59812-7056; 888-268-2743 or 406-243-5467; 406-243-4200 (TTY); 406-243-2349 (Fax); [rtcrural@mso.umt.edu](mailto:rtcrural@mso.umt.edu); <http://rtc.ruralinstitute.umt.edu>

© 2014 RTC:Rural. Our research is supported by grant #H133G110077 from the National Institute on Disability and Rehabilitation Research, U.S. Dept. of Education. The opinions expressed reflect those of the authors and are not necessarily those of the funding agency.

