

University of Montana

ScholarWorks at University of Montana

Chemistry and Biochemistry Faculty
Publications

Chemistry and Biochemistry

1-21-2015

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May

Colorado State University - Fort Collins

T. Lee

Colorado State University - Fort Collins

G. R. McMeeking

Colorado State University - Fort Collins

Sheryl Kashi Akagi

The University of Montana, sheryl.akagi@gmail.com

A. P. Sullivan

Colorado State University - Fort Collins

See next page for additional authors

Follow this and additional works at: https://scholarworks.umt.edu/chem_pubs



Part of the [Biochemistry Commons](#), and the [Chemistry Commons](#)

Let us know how access to this document benefits you.

Recommended Citation

May, A. A.; Lee, T.; McMeeking, G. R.; Akagi, Sheryl Kashi; Sullivan, A. P.; Urbanski, S. P.; Yokelson, Robert; and Kreidenweis, S. M., "Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes" (2015). *Chemistry and Biochemistry Faculty Publications*. 90.
https://scholarworks.umt.edu/chem_pubs/90

This Article is brought to you for free and open access by the Chemistry and Biochemistry at ScholarWorks at University of Montana. It has been accepted for inclusion in Chemistry and Biochemistry Faculty Publications by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact scholarworks@mso.umt.edu.

Authors

A. A. May, T. Lee, G. R. McMeeking, Sheryl Kashi Akagi, A. P. Sullivan, S. P. Urbanski, Robert Yokelson, and S. M. Kreidenweis



Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May^{1,*}, T. Lee^{1,**}, G. R. McMeeking^{1,***}, S. Akagi², A. P. Sullivan¹, S. Urbanski³, R. J. Yokelson², and S. M. Kreidenweis¹

¹Department of Atmospheric Science, Colorado State University, Fort Collins, CO, USA

²Department of Chemistry, University of Montana, Missoula, MT, USA

³Missoula Fire Sciences Laboratory, Rocky Mountain Research Station, US Forest Service, Missoula, MT, USA

* now at: Department of Civil, Environmental and Geodetic Engineering, The Ohio State University, Columbus, OH, USA

** now at: Department of Environmental Science, Hankuk University of Foreign Studies, Yongin, Korea

*** now at: Droplet Measurement Technologies, Boulder, CO, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Received: 12 December 2014 – Accepted: 5 January 2015 – Published: 21 January 2015

Correspondence to: A. A. May (may.561@osu.edu) and
S. M. Kreidenweis (sonia@atmos.colostate.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

15, 1953–1988, 2015

**Observations and
analysis of organic
aerosol evolution in
some prescribed fire
smoke plumes**

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Open biomass burning is a significant source of primary air pollutants such as particulate matter and non-methane organic gases. However, the physical and chemical atmospheric processing of these emissions during transport is poorly understood. Atmospheric transformations of biomass burning emissions have been investigated in environmental chambers, but there have been limited opportunities to investigate these transformations in the atmosphere. In this study, we deployed a suite of real-time instrumentation on a Twin Otter aircraft to sample smoke from prescribed fires in South Carolina, conducting measurements at both the source and downwind to characterize smoke evolution with atmospheric aging. Organic aerosol (OA) within the smoke plumes was quantified using an Aerosol Mass Spectrometer (AMS), along with refractory black carbon (rBC) using a Single Particle Soot Photometer and carbon monoxide (CO) and carbon dioxide (CO₂) using a Cavity Ring-Down Spectrometer. During the two fires for which we were able to obtain aerosol aging data, normalized excess mixing ratios and “export factors” of conserved species (rBC, CO, CO₂) were unchanged with increasing sample age. Investigation of AMS mass fragments indicated that the in-plume fractional contribution ($f_{m/z}$) to OA of the primary fragment (m/z 60) decreased downwind, while the fractional contribution of the secondary fragment (m/z 44) increased. Increases in f_{44} are typically interpreted as indicating chemical production of secondary OA (SOA). Likewise, we observed an increase in the O:C elemental ratio downwind, which is usually associated with aerosol aging. However, the rapid mixing of these plumes into the background air suggests that these chemical transformations may be attributable to the different volatilities of the compounds that fragment to these m/z in the AMS. The gas-particle partitioning behavior of the bulk OA observed during the study was consistent with the predictions from a parameterization developed for open biomass burning emissions in the laboratory. Furthermore, we observed no statistically-significant increase in total organic mass with atmospheric transport. Hence, our results suggest that dilution-driven evaporation likely dominated over

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



these transformations to specific physical and chemical processes is difficult (Heilman et al., 2014).

Laboratory studies have been conducted to attempt to separate these processes for biomass-burning-derived OA. As part of the third Fire Lab at Missoula Experiment (FLAME-III), May et al. (2013) derived a volatility distribution and related thermodynamic parameters representative of the primary emissions from all of the biomass fuels studied. In that same study, Hennigan et al. (2011) and Ortega et al. (2013) investigated chemical transformations of the emissions using an environmental chamber and a potential aerosol mass chamber, respectively. Results from both chambers demonstrated that the OA mass can be enhanced, depleted, or remain roughly constant with oxidation, similar to field measurements, yet the OA became apparently more oxidized with photochemical aging, as interpreted from the organic mass fragments measured via online aerosol mass spectrometry.

In this work, we report and interpret observations from the South Carolina fire Emissions And Measurements (SCREAM) campaign conducted in October-November 2011 (Akagi et al., 2013, 2014; May et al., 2014; Sullivan et al., 2014). The objectives of SCREAM were to simulate moderately intense wildfires by conducting prescribed burns at sites with high fuel loadings, to characterize the emissions and develop estimates of emission factors and emission ratios from both ground- and aircraft-based sampling, and to sample plumes downwind as they evolved during atmospheric transport. We also sampled fires of opportunity during the study. The SCREAM campaign was the first study, to our knowledge, to include simultaneous aircraft-based online measurements of refractory black carbon (rBC), time-resolved non-refractory sub-micron PM measurements (including OA), and time-resolved water-soluble organic carbon (WSOC) and levoglucosan (LEV) measurements, in addition to a suite of gas-phase compounds. Companion papers have reported airborne trace gas emissions (Akagi et al., 2013), ground-based trace gas emissions (Akagi et al., 2014), airborne WSOC and smoke marker emissions (Sullivan et al., 2014), and airborne primary PM emissions (May et al., 2014). This paper focuses on airborne observations of the OA mass

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentrations and composition near the source and transformations to OA mass concentration and composition during the first hours of atmospheric transport.

2 Methods

Emissions from five of the seven fires sampled during SCREAM are discussed in this paper. Details including fuel type, area burned, meteorology and stand history were provided by Akagi et al. (2013) and are summarized briefly here. Two of the burns were conducted on the Fort Jackson (FJ) Army Base (located northeast of Columbia, SC) in Blocks 9b (FJ 9b; 34°0'15" N, 80°52'37" W; 1 November 2011) and 22b (FJ 22b; 34°5'4" N, 80°52'16" W; 2 November 2011). These burns occurred in older stands that had not been treated for a number of years, and were intended to simulate wildfires. Fuel inventories indicated that vegetation comprised primarily mature longleaf pine (*Pinus palustris*) and loblolly pine (*Pinus taeda*) with some contributions of turkey oak (*Quercus laevis Walter*) and farkleberry (*Vaccinium arboretum Marsh.*). Complementary ground-based measurements of emissions from the FJ burns were reported by Akagi et al. (2014). The three other sampled fires were designated Georgetown (33°12'9" N, 79°24'6" W; 7 November 2011), Francis Marion (33°12'55" N, 79°28'34" W; 8 November 2011), and Bamberg (33°14'5" N, 80°56'41" W; 10 November 2011), based on the location in SC where the fire occurred. Georgetown and Francis Marion were located in coastal SC, likely burning coastal grasses and longleaf pine understory, respectively, based on in-flight observations. The Bamberg fire, located roughly 80 km due south of the Fort Jackson site in inland SC, was likely comprised of multiple fuel types, including longleaf/loblolly pine understory as well as marsh grasses, based on smoke marker ratio measurements reported in Sullivan et al. (2014).

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was conducted from a low turbulence inlet (Wilson et al., 2004) followed by a non-rotating Micro Orifice Uniform Deposit Impactor (MOUDI; Marple et al., 1991). The MOUDI was operated such that it served as a PM_{10} impactor (i.e., having 50 % particle transmission efficiency for particulate matter of 1 μm aerodynamic diameter with a sharpness of 1.08 – particles less than roughly 900 nm will be transmitted with 100 % efficiency). For consistency, all data were adjusted to the same timestamp, which we defined from HR-ToF-AMS.

2.2.1 Aerosol mass spectrometer

The HR-ToF-AMS (hereafter AMS) characterizes non-refractory sub-micron aerosol by focusing sampled particles through an aerodynamic lens, collecting particles on a thermal vaporizer, ionizing the vaporized particles via electron impactation, and detecting ions (m/z) in the high-resolution time-of-flight mass spectrometer (DeCarlo et al., 2006). Using the ToF-AMS data analysis toolkit SQUIRREL 1.53G/PIKA 1.12G (Sueper et al., 2013), aerosol mass concentrations can be reconstructed from the m/z signal; for this study, we fit HR peaks for $m/z \leq 200$. These concentrations are dependent on instrument parameters (e.g., ionization efficiency and vaporizer collection efficiency). Ionization efficiency calibrations were performed with 350 nm ammonium nitrate particles throughout the campaign, with values ranging from 1.83×10^{-7} to 2.91×10^{-7} . Composition-dependent collection efficiencies were calculated following the algorithm of Middlebrook et al. (2012), which is now built into the SQUIRREL software, for each AMS sample and ranged from roughly 0.5–0.9, with a campaign-average value of 0.53. We report AMS-derived emissions data of nitrate, sulfate, ammonium, and chloride elsewhere (May et al., 2014).

The AMS was mounted into National Center for Atmospheric Research GV-type aircraft racks with a pressure-controlled inlet to reduce fluctuations of the pressure within the aerodynamic lens (Bahreini et al., 2008). During operation, data were exclusively collected using the “V-mode” of the ion time-of-flight within the mass spectrometer; no

particle time-of-flight data were collected. AMS data were typically collected with a time resolution of 6 s (corresponding to a distance of roughly 250–300 m).

While we obtained simultaneous measurements of gas-phase CO₂, we have not corrected our data for any potential interference with the signal at *m/z* 44 (CO₂⁺) in the AMS. The AMS samples particles roughly 10⁷ times more efficiently than the gas-phase. We estimate that on average, our plume OA concentrations are positively biased by 0.0044 ± 0.0019 % (both near the source and downwind), our background OA concentrations are positively biased by 0.025 ± 0.021 %, and our *m/z* 44 measurements are positively biased by 0.20 ± 0.11 %, all based on co-located gas-phase CO₂ measurements. Consequently, we deemed this correction unnecessary as this interference represents < 0.5 % of our reported values.

2.2.2 Single particle soot photometer

The SP2 provides operationally-defined rBC mass concentrations via laser-induced incandescence (Stephens et al., 2003; Schwarz et al., 2006). Absorbing material present in particles is heated to its vaporization temperature and emits radiation, which is measured by optical detectors. This approach removes uncertainties due to interferences of artifacts that have been observed during filter-based approaches (Kirchstetter et al., 2004) and excludes the influence of “brown” carbon that can bias optical absorption methods (Andreae and Gelencsér, 2006; Lack et al., 2012), although it has been shown the method responds to some metals. Signal is related to rBC mass via calibration procedures; during SCREAM, calibrations were performed using fullerene soot. Generally, rBC mass concentrations were recorded every 6 s, similar to the AMS. Additional details related to the SP2 operation during this campaign can be found in May et al. (2014).

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2.3 Cavity ring-down spectrometer

A Picarro G2401 provided 1 Hz measurements of CO₂, CO, CH₄, and H₂O, which are the major gas-phase emissions from combustion sources. The CRDS was calibrated in-flight using mixed standards of CO₂/CO/CH₄ and procedures similar to those described by Urbanski (2013). These data were applied to calculate emission ratios and emission factors of particle-phase species, as described below.

2.2.4 Airborne fourier-transform infrared spectrometer

The AFTIR data used and obtained during this study are described in detail by Akagi et al. (2013). “Grab samples” were collected in background air and as the Twin Otter traversed the smoke plume; these samples were held for 2–3 min of signal-averaging in order to improve sensitivity. Compound mixing ratios were quantified via a non-linear least-squares method for the majority of the identifiable trace gases (Burling et al., 2011; Yokelson et al., 2007). These AFTIR data provide complementary results describing gas-phase chemistry that assist the interpretation of particle-phase measurements.

2.3 Data processing

2.3.1 Plume identification and age

Emissions were rapidly diluted and mixed within the boundary layer (within roughly 30 min downwind), and the plumes did not penetrate into the free troposphere, so visual tracking of the plumes was challenging. Furthermore, the plume from the FJ 22b fire entered restricted air space near Columbia, SC, so it was only possible to follow this plume for a short distance from the point of emission. Positive downwind plume intercepts were confirmed through simultaneous spikes in measured values of OA, BC, and CO. These confirmed intercepts were corroborated by simultaneous increases in smoke markers (Sullivan et al., 2014) and trace gases (Akagi et al., 2013). Intercept-

1962

ACPD

15, 1953–1988, 2015

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



averaged concentrations for each transect were derived by integrating the excess area (above background) under the data time series curves and dividing by the elapsed time in the window of integration.

Sample ages (times since emission) were estimated using the distance from the source and mean wind speed. Distance from the source was computed using the haversine formula and the spatial coordinates measured by the AIMMS-20. Mean wind speed was also measured using the AIMMS-20. Akagi et al. (2013) estimated that this approach has an uncertainty of roughly 30 %, largely due to uncertainties in the wind speed data. Due to the nature of the flight path, plume intercepts rarely occurred perpendicular to the plume; in fact, they were often diagonal transects. Thus, a given sample can be associated with a range of estimate ages. In subsequent figures, we plot the average age of a plume intercept along with error bars representing the range of ages; in these figures, we do not include the estimated uncertainty of 30 % on this range.

This calculation provides a pseudo-Lagrangian estimate of plume age, and hence, a downwind sample can be related to emissions measured directly at the source at the appropriate earlier time. Therefore, some of the data we present are labeled “Lagrangian” when we had corresponding source measurements, whereas for others (labeled “non-Lagrangian”) we can estimate time since emission based on distance from the source and ambient wind speed but do not have a corresponding source sample. For consistency with May et al. (2014), we defined “near-source” samples as those collected within 5 km of the fire, while downwind samples were those collected at distances greater than 5 km.

In-plume data from all research flights were corrected for local background concentrations via integration under the curves in data time series between out-of-plume measurements. The resulting species concentrations are “excess” concentrations and denoted by the delta symbol, i.e., ΔX is the excess concentration of species X . We show background values of some parameters in some of the following figures; these back-

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ground concentrations represent the median background concentration for the duration of the given flight. Sample background-corrected data are provided in Fig. 1.

2.3.2 Excess emission ratios and emission factors

Normalized excess mixing ratios (NEMR) are often used to account for transient fire behavior and the dilution and mixing of plumes with background air during transport (e.g., Hobbs et al., 2003) and are defined as:

$$\text{NEMR}_X = \frac{\Delta X}{\Delta \text{CO}} \quad (1)$$

where ΔX is the excess concentration of species X , and ΔCO is the background-corrected value of CO. Since both numerator and denominator are excess quantities, uncertainties in their values increase as the plume dilutes and in-plume concentrations approach the background concentrations. Here, we report instantaneous plume-integrated NEMR_X for each plume interception, so our values will differ from the “fire-integrated” values reported in May et al. (2014). NEMR_X are reported here in units of $\mu\text{g m}^{-3} (\text{ppmv CO})^{-1}$; this value can be converted to $\text{g}(\text{g CO})^{-1}$ by multiplying by a factor of $8.7 \times 10^{-4} \text{ ppmv CO} (\mu\text{g CO m}^{-3})^{-1}$. Strictly speaking, NEMR_X is a misnomer for aerosol mass concentrations, but we utilize this terminology for consistency with the vast body of prior literature.

Time series of instantaneous NEMR_X provide information on transient smoke behavior (Jolleys et al., 2014). By associating instantaneous NEMR_X with time since emission, physicochemical transformations can be investigated, since NEMR_X accounts for dilution and thus should be constant with time in the absence of sources or sinks of the species X . The net formation of secondary organic aerosol (SOA) in smoke plumes can be inferred from an increase in NEMR_{OA} with distance downwind (Yokelson et al., 2009; Akagi et al., 2012). On the other hand, since OA emitted from biomass burning sources is semi-volatile, net evaporation of particle-phase mass as dilution proceeds would appear as a decrease in NEMR_{OA} .

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Emission factors (EF) are widely used descriptors of fire emissions (Ward and Radke, 1993; Andreae and Merlet, 2001). Their calculation relates the mass of X emitted (M_X) to the mass of dry fuel consumed (M_{fuel}). In cases where the mass of fuel consumed is unknown, a carbon mass balance approach can be applied, which relates the change in the concentration of X relative to the background (ΔX ; $\mu\text{g m}^{-3}$) to the excess carbon concentrations (i.e., background-corrected concentrations that have been converted to mg-C m^{-3}) of CO_2 , CO , total organic gases ($\text{TOG} = \text{CH}_4 + \text{NMOG}$), and carbonaceous PM (PM_C):

$$\text{EF}_X = \frac{M_X}{M_{\text{fuel}}} = \frac{\Delta X}{\Delta\text{CO}_2 + \Delta\text{CO} + \Delta\text{TOG} + \Delta\text{PM}_C} f_C \quad (2)$$

In Eq. (2), the term f_C is a conversion factor representing fuel carbon content. Since we lack detailed fuel information, we assume that $f_C = 0.50$, roughly the average fuel carbon content of southeastern (SE) US coastal plain biomass fuels reported in laboratory studies (Burling et al., 2010; May et al., 2014; McMeeking et al., 2009); we also lack an estimate of TOG in the emissions. Furthermore, $\Delta\text{PM}_C \ll (\Delta\text{CO}_2 + \Delta\text{CO})$. Hence, we approximate EF_X neglecting both ΔTOG and ΔPM_C , which may result in an over-estimate in EF_X of 3–4% (Yokelson et al., 2013). Like NEMR_X , EF_X are based on excess concentrations and account for dilution, but if an “emission factor” is computed with downwind data, the value obtained reflects any sources or sinks of the originally-emitted species X . Hereafter, we will refer to downwind “emission factors” as “export factors”, also denoted as EF_X and calculated from Eq. (2); the main distinction is that an export factor describes X downwind from the source, and thus may be subject to atmospheric transformations. We report EF_X as $\text{mg}(\text{kg dry fuel})^{-1}$.

3 Results and discussion

During the study, only two fires provided adequate downwind aerosol data allowing us to investigate in-plume aerosol physicochemical transformations: the FJ 9b fire and

1965

ACPD

15, 1953–1988, 2015

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the Francis Marion fire. As mentioned earlier, the plume from the FJ 22b fire entered restricted airspace and could not be pursued. Further, the Georgetown fire was a small fire whose plume rapidly mixed with the background, so downwind ΔOA was small and uncertain; the Bamberg samples represented two distinct fuel types as shown elsewhere (May et al., 2014; Sullivan et al., 2014), making it difficult to distinguish transformations during transport from differences in the sources.

In Fig. 2, we present composition data vs. estimated time since emission of the NEMR or EF of four major components present in the biomass burning smoke sampled for the FJ 9b fire: OA (NEMR_{OA} ; Fig. 2a), rBC (NEMR_{rBC} ; Fig. 2b), CO (EF_{CO} ; Fig. 2c), and CO_2 (EF_{CO_2} ; Fig. 2d). Data near the source are presented as box-and-whisker plots (25th–75th and 10th–90th percentiles); these data were collected over roughly 2.5 h of real-time, during which the modified combustion efficiency (MCE) (Ward and Radke, 1993) varied between 0.900 and 0.930, which explains some of the variability in the data. Data up to five hours downwind were obtained and are shown as closed symbols for Lagrangian points, or open symbols for non-Lagrangian data. For downwind samples, vertical errors bars represent estimated measurement uncertainties while horizontal error bars represent the range of estimated plume ages for non-perpendicular plume transects; horizontal error bars do not account for the estimated 30 % measurement uncertainty in wind speed.

We expect rBC, CO, and CO_2 to be conserved with transport since they are stable in the atmosphere on the timescales considered here. Indeed, unpaired t tests for the data shown in Fig. 2b–d indicate that there was no significant difference between these species at the source and downwind (two-tailed p values > 0.13). Differences in mean downwind EFs are attributable to measurement uncertainties, including identification of the plume edges, and variability in the combustion phase at the source. Based on the exponential decay with distance from the source of absolute mixing ratios of CO and CO_2 , we infer an average mixing rate (the inverse of the dilution timescale) of 1.6 h^{-1} during the FJ 9b experiment.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Since OA is reactive and semi-volatile, it is perhaps not surprising that the downwind $NEMR_{OA}$ over 2–5 h of atmospheric aging is significantly lower than the $NEMR_{OA}$ at the source (Fig. 2a; two-tailed p value = 0.015), suggesting a net loss of emitted OA via evaporation and/or reaction. As demonstrated by Akagi et al. (2013), the smoke plume was photochemically active, as evident through enhancements of ozone and formaldehyde relative to the source.

Figure 3 is identical to Fig. 2 but represents the Francis Marion burn, the only other case with adequate downwind aerosol measurements (here, up to 1.5 h after emission) and no other known biomass burning emission sources. Akagi et al. (2013) inferred photochemical processing was occurring in the Francis Marion plume, based on observed downwind enhancements of ozone and formaldehyde relative to the source. However, unlike the FJ 9b fire, none of the computed downwind NEMR and EF shown in Fig. 3 were significantly different from the source (all two-tailed p values > 0.32). The background OA concentrations, which we assume contribute to gas-particle partitioning of emitted OA by providing additional absorptive material, were roughly 50 % greater during the Francis Marion fire compared to the FJ 9b fire; furthermore, the dilution rate was 20 % slower for the Francis Marion plume (1.3 h^{-1}), and the plume was observed over a much shorter time period. These factors would slow the evaporation of emitted OA, and limit the time over which chemical transformations could occur and be observed. Indeed, over the first 1.5 h after emission, the data for FJ 9b shown in Fig. 2 also indicated no statistically-significant detectable change in $NEMR_{OA}$.

3.1 Chemical transformations of organic aerosol

In this section, we utilize two approaches to investigate chemical transformations of the organic aerosol for the two cases with downwind data, the FJ 9b and Francis Marion burns: fragment evolution (Fig. 4) and elemental ratio analyses (Fig. 5). Ng et al. (2010) demonstrated that “fresh” OA in ambient samples can be distinguished by organic fragment signatures in the mass spectra (e.g., $C_3H_7^+$ at m/z 43), while “aged” OA is more highly oxidized and can be distinguished by a strong contribution of CO_2^+ (m/z 44).

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The fractional contributions of each of these fragments to the total OA concentration (e.g., $f_{44} = C_{44}/C_{OA}$, where C_{44} is the mass concentration of particulate CO_2^+) change with atmospheric aging: f_{43} is expected to decrease and f_{44} to increase. Levoglucosan and other anhydrosugars that are combustion products of cellulose, and thus used as molecular markers for biomass burning emissions (Simoneit et al., 1999; Sullivan et al., 2008), can be identified in AMS spectra at m/z 60 ($C_2H_4O_2^+$) (Alfarra et al., 2007; Lee et al., 2010). May et al. (2012) and references therein demonstrated that levoglucosan is semi-volatile at ambient conditions; hence, m/z 60 is expected to evaporate during dilution, if this finding is extrapolated to all anhydrosugars in general. Furthermore, Hennigan et al. (2010) demonstrated that levoglucosan is reactive and chemically decays similar to the hydrocarbon-like (m/z 43) fragments. Cubison et al. (2011) and Ortega et al. (2013) thus modified the Ng et al. (2010) approach, comparing f_{60} and f_{44} to infer photochemical aging of BBOA.

In Fig. 4, we present excess f_{60} (Δf_{60}) and excess f_{44} (Δf_{44}) for the FJ 9b and Francis Marion fires. These excess fragment fractional contributions were computed from background-corrected m/z 60 or m/z 44 mass concentrations by dividing that excess concentration by ΔOA . Thus, as the plume dilutes and becomes less distinguishable from the background, Δf_{60} and Δf_{44} should remain constant if neither preferentially evaporates, reacts, or accumulates within the plume. For the FJ 9b fire, the source-downwind differences for both Δf_{60} (Fig. 4a) and Δf_{44} (Fig. 4b) are statistically significant (two-tailed p value < 0.0001). For the Francis Marion fire, Δf_{60} (Fig. 4c) is significantly lower at the source than downwind (two-tailed p value < 0.0001), while Δf_{44} (Fig. 4d) is significantly higher downwind than at the source (two-tailed p value = 0.029). The result for Δf_{60} is consistent with Fig. 2a; that is, the decrease in Δf_{60} downwind during the FJ 9b fire reflects the decrease in $NEMR_{OA}$. An observed decrease in Δf_{60} with no decrease in OA concentration during the Francis Marion fire may be related to chemical reactions of compounds that fragment to m/z 60 or to differences in the volatility of these compounds compared to the bulk OA. The mechanistic driver of all transformations will be explored below.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The increase in Δf_{44} with plume age for both fires indicates a compositional change toward increasing mass fractional contributions from molecules that fragment to CO_2^+ . If only dilution (and hence, evaporation) was occurring in the plumes as they moved downwind, Δf_{44} should be conserved, provided its parent's volatility is similar to that of the bulk of the emitted OA. The observed increase in CO_2^+ in these photochemically-active environments may indicate that production of SOA occurred within the plumes, although there were no statistically-significant increases in the measured downwind NEMR_{OA} , as also found in some previous field studies (e.g., Capes et al., 2008; Cubison et al., 2011). On the other hand, this increase could also indicate that the species fragmenting to m/z 44 are relatively less volatile than the bulk OA that evaporates during transport and dilution.

There is experimental evidence investigating chemically-resolved volatility that is consistent with the evaporation of bulk OA resulting in a relative increase in m/z 44 and a relative decrease in m/z 60. Huffman et al. (2009a) demonstrated for ambient samples in two different megacities that, at a given temperature in a thermodenuder, m/z 60 evaporated to a greater extent than the bulk OA, while m/z 44 evaporated to a lesser extent than the bulk OA. While heating OA is technically not the same as diluting OA, the response of OA to increased temperature is analogous to the response of OA to increased dilution. Furthermore, Collier and Zhang (2013) demonstrated that f_{44} increased with decreasing C_{OA} for vehicle test data in the absence of chemistry and hypothesized that this observation was attributable to preferential evaporation of less-oxidized OA species. Thus, the observed changes during SCREAM in Δf_{44} and Δf_{60} may be due, at least in part, to physical changes occurring as some of the emitted OA is volatilized upon dilution with ambient air.

Another framework for tracking the chemical evolution of OA was suggested by Heald et al. (2010), who proposed the use of elemental ratios (hydrogen to carbon, H:C, and oxygen to carbon, O:C) to describe photochemical aging of OA. Similar to the fragment evolution, with increasing OH exposure, H:C is expected to decrease (e.g., due to hydrogen abstraction reactions) and O:C is expected to increase (e.g., due to

oxygen addition to alkyl radicals). In Fig. 5 we present the evolution of the elemental ratio of H:C and O:C during atmospheric transport of the biomass burning plumes from both fires; values of the ratios for the average background, out-of-plume ratio for each fire are shown as dotted lines. We assume uncertainties of 31 % of O:C and 10 % for H:C, based on Aiken et al. (2008).

For both fires, the average background H:C ratio was roughly 15 % greater than the H:C at the source; downwind H:C values were mostly within the source variability. As the plumes were transported downwind and mixed with background OA, based on measured dilution rates we expected H:C to have increased toward the background values on a 2 to 3 h timescale if it were a conserved tracer. The lack of a clear decrease with time since emission in both experiments suggests either that loss of both H and C occurred in the plume, or increases in C occurred without corresponding addition of H that would maintain the H:C observed at the source. For O:C, about half the downwind values were higher than could be explained by measured variability at the source, and the background OA had O:C within (but at the lower end) of the range at the source. Dilution with background air was thus expected to have had little impact on O:C if O:C were a conserved tracer. Like m/z 44, O:C could have increased with time if photochemical production and condensation of high O:C species occurred, or potentially by a solely physical process if C were lost from the aerosol phase by reaction or evaporation of species that had lower O:C than the average observed at the source; in fact, Huffman et al. (2009b) demonstrated that O:C increased and H:C decreased with increasing evaporation of bulk OA in biomass burning emissions during thermodenuder experiments.

3.2 Physical transformations of organic aerosol

A net loss of OA due solely to dilution-driven evaporation may thus be consistent with the observations in Figs. 2–5. However, we note that we cannot definitively state that no aerosol chemistry has occurred within the plumes as they age. In the following, we assume a priori knowledge that dilution-driven evaporation dominates over chemical

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



processing and apply results obtained in prior lab studies of biomass burning emissions to simulate plume evolution to test this assumption.

Simulations representing the process of dilution alone are presented in Fig. 6, which shows EF_{OA} data (representing the emission factors near the source and export factors downwind) as a function of the total mass concentration of observed organic aerosol (i.e., not background corrected), C_{OA} , for six flights. Model curves were calculated using the following equation (Donahue et al., 2006; Robinson et al., 2010):

$$EF_{OA} = EF_{tot} \sum_i f_i \left(1 + \frac{C_i^*}{C_{OA}} \right)^{-1} \quad (3)$$

where i represents arbitrarily-chosen surrogate compounds defined by their saturation concentration (C_i^* ; related to saturation vapor pressure through the ideal gas law), and f_i is the mass fraction of each species i relative to the total emitted organics. The set of f_i and C_i^* is referred to as a volatility distribution. Here, we utilize the volatility distribution for emissions from open biomass burning that was proposed by May et al. (2013), which is comprised of surrogate compounds representing seven logarithmically-spaced C_i^* bins. C_{OA} represents the total OA concentration (emissions + background).

EF_{tot} is the emission factor of total organics (gas + particle phase) that are constrained by the volatility distribution (here, all material between $C_i^* = 3 \times 10^{-3} \mu\text{g m}^{-3}$ and $3 \times 10^4 \mu\text{g m}^{-3}$, so this is not equivalent to NMOG), and hence, contribute to gas-particle partitioning; EF_{tot} is likely dominated by biomass-burning-derived organics but may include background semi-volatile organic material that can partition into the particle phase due to the presence of the biomass burning smoke. Values of EF_{tot} were inferred using Eq. (3) with measured C_{OA} , calculated EF_{OA} (from Eq. 2), and the volatility distribution from May et al. (2013) as inputs for each plume intercept. In Fig. 6, the lines represent predictions based on the average EF_{tot} inferred for each fire, while the shaded areas represent \pm one SD in EF_{tot} . Eq. (3) implies that EF_{OA} (regardless of whether this represents an emission factor or export factor) decreases with increasing dilution, due to the physical repartitioning of semi-volatile species.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are represented using a single set of model inputs, which does not account for this variability with MCE. While other factors likely play a role, these two are likely to be the most important.

4 Conclusions

In this work, we present field observations of the physicochemical evolution of the organic aerosol present in biomass burning plumes from two prescribed fires in South Carolina. Downwind observations of rBC to CO ratios and emission factors of CO, and CO₂ are not statistically different from those at the source. The ratio of OA to CO was significantly lower than at the source during the fire that we were able to follow downwind for up to five hours of atmospheric aging, but this ratio was not significantly different downwind for the other fire, which may be related to the much shorter observable atmospheric aging time (~ 2 h).

We observed significant differences in downwind ratios of AMS mass fragments thought to be indicative of fresh biomass burning emissions (*m/z* 60, which decreased) and more oxidized OA species (*m/z* 44, which increased), consistent with prior reported laboratory photo-oxidation experiments. While the observed increases in Δf_{44} (and the O:C ratio) imply the possibility of SOA production within the plume, these observed changes are also consistent with differences in the volatilities of the species fragmenting to *m/z* 60 and *m/z* 44 relative to the bulk OA, resulting in differences in evaporation as the plume dilutes into background air.

Our observations and model simulations suggest that dilution-driven evaporation out of the particle phase dominates over condensation of semi-volatile material into the particle phase over roughly the first hour of transport, after which the OA in the plume reached an equilibrium state with the background in our observations. This finding is consistent with results from previous literature (Akagi et al., 2012; Jolleys et al., 2012); however, other studies report increases in OA with increasing plume age (DeCarlo et al., 2008; Vakkari et al., 2014; Yokelson et al., 2009). The exact cause of this variability

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ity in observations is unclear. These remaining unexplained differences among different field studies highlight the need for additional research on atmospheric physicochemical transformations of biomass burning plumes.

Acknowledgements. We acknowledge funding from the Joint Fire Science Program under project JFSP 11-1-5-12 to S. M. Kreidenweis and the Strategic Environmental Research and Development Program project RC-1649 administered partly through the Forest Service Research Joint Venture Agreement 08JV11272166039 to R. Yokelson. Additional flight hours and CRDS data were provided by Joint Fire Science Program project 08-1-6-09 to S. Urbanski. Adaptation of the Twin Otter was supported by NSF grants ATM-0531044 and ATM-0936321 to R. Yokelson. We also thank Ezra Levin and the NSF/NCAR Research Aviation Facility for assistance with the installation of instruments to the Twin Otter; Shane Murphy, Roya Bahreini, and Ann Middlebrook for guidance in modifying the CSU AMS for aircraft operation; and Jose Jimenez, Tim Onasch, Jill Craven, and Misha Schurman for discussions related to AMS data analysis. This work would not have been possible without the support of the Twin Otter Science Team, especially pilot Bill Mank and mechanic Steve Woods; John Maitland and the forestry staff at Fort Jackson who conducted those prescribed fires; and the Columbia Dispatch Office of the South Carolina Forestry Commission for providing additional prescribed fire locations during the campaign.

References

- Aiken, A. C., Decarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, 42, 4478–4485, doi:10.1021/es703009q, 2008.
- Akagi, S. K., Yokelson, R. J., Wiedinmyer, C., Alvarado, M. J., Reid, J. S., Karl, T., Crounse, J. D., and Wennberg, P. O.: Emission factors for open and domestic biomass burning for use in atmospheric models, *Atmos. Chem. Phys.*, 11, 4039–4072, doi:10.5194/acp-11-4039-2011, 2011.

ACPD

15, 1953–1988, 2015

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Akagi, S. K., Craven, J. S., Taylor, J. W., McMeeking, G. R., Yokelson, R. J., Burling, I. R., Urbanski, S. P., Wold, C. E., Seinfeld, J. H., Coe, H., Alvarado, M. J., and Weise, D. R.: Evolution of trace gases and particles emitted by a chaparral fire in California, *Atmos. Chem. Phys.*, 12, 1397–1421, doi:10.5194/acp-12-1397-2012, 2012.
- 5 Akagi, S. K., Yokelson, R. J., Burling, I. R., Meinardi, S., Simpson, I., Blake, D. R., McMeeking, G. R., Sullivan, A., Lee, T., Kreidenweis, S., Urbanski, S., Reardon, J., Griffith, D. W. T., Johnson, T. J., and Weise, D. R.: Measurements of reactive trace gases and variable O₃ formation rates in some South Carolina biomass burning plumes, *Atmos. Chem. Phys.*, 13, 1141–1165, doi:10.5194/acp-13-1141-2013, 2013.
- 10 Akagi, S. K., Burling, I. R., Mendoza, A., Johnson, T. J., Cameron, M., Griffith, D. W. T., Paton-Walsh, C., Weise, D. R., Reardon, J., and Yokelson, R. J.: Field measurements of trace gases emitted by prescribed fires in southeastern US pine forests using an open-path FTIR system, *Atmos. Chem. Phys.*, 14, 199–215, doi:10.5194/acp-14-199-2014, 2014.
- 15 Alfara, M. R., Prévôt, A. S. H., Szidat, S., Sandradewi, J., Weimer, S., Lanz, V. A., Schreiber, D., Mohr, M., and Baltensperger, U.: Identification of the mass spectral signature of organic aerosols from wood burning emissions, *Environ. Sci. Technol.*, 41, 5770–5777, doi:10.1021/es062289b, 2007.
- Andrae, M. O. and Gelencsér, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 6, 3131–3148, doi:10.5194/acp-6-3131-2006, 2006.
- 20 Andrae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, doi:10.1029/2000GB001382, 2001.
- Bahreini, R., Dunlea, E. J., Matthew, B. M., Simons, C., Docherty, K. S., DeCarlo, P. F., Jimenez, J. L., Brock, C. A., and Middlebrook, A. M.: Design and operation of a pressure-controlled inlet for airborne sampling with an aerodynamic aerosol lens, *Aerosol Sci. Tech.*, 25, 42, 465–471, doi:10.1080/02786820802178514, 2008.
- 30 Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: a scientific assessment, *J. Geophys. Res.-Atmos.*, 118, 5380–5552, doi:10.1002/jgrd.50171, 2013.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Burling, I. R., Yokelson, R. J., Griffith, D. W. T., Johnson, T. J., Veres, P., Roberts, J. M., Warneke, C., Urbanski, S. P., Reardon, J., Weise, D. R., Hao, W. M., and de Gouw, J.: Laboratory measurements of trace gas emissions from biomass burning of fuel types from the southeastern and southwestern United States, *Atmos. Chem. Phys.*, 10, 11115–11130, doi:10.5194/acp-10-11115-2010, 2010.
- Burling, I. R., Yokelson, R. J., Akagi, S. K., Urbanski, S. P., Wold, C. E., Griffith, D. W. T., Johnson, T. J., Reardon, J., and Weise, D. R.: Airborne and ground-based measurements of the trace gases and particles emitted by prescribed fires in the United States, *Atmos. Chem. Phys.*, 11, 12197–12216, doi:10.5194/acp-11-12197-2011, 2011.
- Capes, G., Johnson, B., McFiggans, G., Williams, P. I., Haywood, J., and Coe, H.: Aging of biomass burning aerosols over West Africa: aircraft measurements of chemical composition, microphysical properties, and emission ratios, *J. Geophys. Res.*, 113, D00C15, doi:10.1029/2008JD009845, 2008.
- Christian, T. J., Kleiss, B., Yokelson, R. J., Holzinger, R., Crutzen, P. J., Hao, W. M., Saharjo, B. H., and Ward, D. E.: Comprehensive laboratory measurements of biomass-burning emissions: 1. Emissions from Indonesian, African, and other fuels, *J. Geophys. Res.*, 108, 4719, doi:10.1029/2003JD003704, 2003.
- Collier, S. and Zhang, Q.: Gas-phase CO₂ subtraction for improved measurements of the organic aerosol mass concentration and oxidation degree by an aerosol mass spectrometer., *Environ. Sci. Technol.*, 47, 14324–31, doi:10.1021/es404024h, 2013.
- Cubison, M. J., Ortega, A. M., Hayes, P. L., Farmer, D. K., Day, D., Lechner, M. J., Brune, W. H., Apel, E., Diskin, G. S., Fisher, J. A., Fuelberg, H. E., Hecobian, A., Knapp, D. J., Mikoviny, T., Riemer, D., Sachse, G. W., Sessions, W., Weber, R. J., Weinheimer, A. J., Wisthaler, A., and Jimenez, J. L.: Effects of aging on organic aerosol from open biomass burning smoke in aircraft and laboratory studies, *Atmos. Chem. Phys.*, 11, 12049–12064, doi:10.5194/acp-11-12049-2011, 2011.
- DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer., *Anal. Chem.*, 78, 8281–9, doi:10.1021/ac061249n, 2006.
- DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crouse, J., Wennberg, P. O., Emmons, L., Shinozuka, Y., Clarke, A., Zhou, J., Tomlinson, J., Collins, D. R., Knapp, D., Weinheimer, A. J., Montzka, D. D., Campos, T., and Jimenez, J. L.:

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Fast airborne aerosol size and chemistry measurements above Mexico City and Central Mexico during the MILAGRO campaign, *Atmos. Chem. Phys.*, 8, 4027–4048, doi:10.5194/acp-8-4027-2008, 2008.

Donahue, N. M., Chuang, W., Epstein, S. A., Kroll, J. H., Worsnop, D. R., Robinson, A. L., Adams, P. J., and Pandis, S. N.: Why do organic aerosols exist? Understanding aerosol lifetimes using the two-dimensional volatility basis set, *Environ. Chem.*, 10, 151, doi:10.1071/EN13022, 2013.

Heald, C. L., Kroll, J. H., Jimenez, J. L., Docherty, K. S., DeCarlo, P. F., Aiken, A. C., Chen, Q., Martin, S. T., Farmer, D. K., and Artaxo, P.: A simplified description of the evolution of organic aerosol composition in the atmosphere, *Geophys. Res. Lett.*, 37, L08803, doi:10.1029/2010GL042737, 2010.

Heilman, W. E., Liu, Y., Urbanski, S., Kovalev, V., and Mickler, R.: Wildland fire emissions, carbon, and climate: plume rise, atmospheric transport, and chemistry processes, *For. Ecol. Manage.*, 317, 70–79, doi:10.1016/j.foreco.2013.02.001, 2014.

Hennigan, C. J., Sullivan, A. P., Collett, J. L., and Robinson, A. L.: Levoglucosan stability in biomass burning particles exposed to hydroxyl radicals, *Geophys. Res. Lett.*, 37, L09806, doi:10.1029/2010GL043088, 2010.

Hennigan, C. J., Miracolo, M. A., Engelhart, G. J., May, A. A., Presto, A. A., Lee, T., Sullivan, A. P., McMeeking, G. R., Coe, H., Wold, C. E., Hao, W.-M., Gilman, J. B., Kuster, W. C., de Gouw, J., Schichtel, B. A., Collett Jr., J. L., Kreidenweis, S. M., and Robinson, A. L.: Chemical and physical transformations of organic aerosol from the photo-oxidation of open biomass burning emissions in an environmental chamber, *Atmos. Chem. Phys.*, 11, 7669–7686, doi:10.5194/acp-11-7669-2011, 2011.

Hobbs, P. V., Sinha, P., Yokelson, R. J., Christian, T. J., Blake, D. R., Gao, S., Kirchstetter, T. W., Novakov, T., and Pilewskie, P.: Evolution of gases and particles from a savanna fire in South Africa, *J. Geophys. Res.*, 108, 8485, doi:10.1029/2002JD002352, 2003.

Hosseini, S., Urbanski, S. P., Dixit, P., Qi, L., Burling, I. R., Yokelson, R. J., Johnson, T. J., Shrivastava, M., Jung, H. S., Weise, D. R., Miller, J. W., and Cocker, D. R.: Laboratory characterization of PM emissions from combustion of wildland biomass fuels, *J. Geophys. Res.-Atmos.*, 118, 9914–9929, doi:10.1002/jgrd.50481, 2013.

Huffman, J. A., Docherty, K. S., Aiken, A. C., Cubison, M. J., Ulbrich, I. M., DeCarlo, P. F., Sueper, D., Jayne, J. T., Worsnop, D. R., Ziemann, P. J., and Jimenez, J. L.: Chemically-

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

resolved aerosol volatility measurements from two megacity field studies, *Atmos. Chem. Phys.*, 9, 7161–7182, doi:10.5194/acp-9-7161-2009, 2009a.

Huffman, J. A., Docherty, K. S., Mohr, C., Cubison, M. J., Ulbrich, I. M., Ziemann, P. J., Onasch, T. B., and Jimenez, J. L.: Chemically-resolved volatility measurements of organic aerosol from different sources, *Environ. Sci. Technol.*, 43, 5351–5357, doi:10.1021/es803539d, 2009b.

Jolleys, M. D., Coe, H., McFiggans, G., Capes, G., Allan, J. D., Crosier, J., Williams, P. I., Allen, G., Bower, K. N., Jimenez, J. L., Russell, L. M., Grutter, M., and Baumgardner, D.: Characterizing the aging of biomass burning organic aerosol by use of mixing ratios: a meta-analysis of four regions., *Environ. Sci. Technol.*, 46, 13093–13102, doi:10.1021/es302386v, 2012.

Jolleys, M. D., Coe, H., McFiggans, G., McMeeking, G. R., Lee, T., Kreidenweis, S. M., Collett, J. L., and Sullivan, A. P.: Organic aerosol emission ratios from the laboratory combustion of biomass fuels, *J. Geophys. Res.-Atmos.*, 119, 12850–12871, doi:10.1002/2014JD021589, 2014.

Kirchstetter, T. W., Novakov, T., and Hobbs, P. V.: Evidence that the spectral dependence of light absorption by aerosols is affected by organic carbon, *J. Geophys. Res.*, 109, D21208, doi:10.1029/2004JD004999, 2004.

Lack, D. A., Langridge, J. M., Bahreini, R., Cappa, C. D., Middlebrook, A. M., and Schwarz, J. P.: Brown carbon and internal mixing in biomass burning particles, *Proc. Natl. Acad. Sci. USA*, 109, 14802–14807, doi:10.1073/pnas.1206575109, 2012.

Lee, T., Sullivan, A. P., Mack, L., Jimenez, J. L., Kreidenweis, S. M., Onasch, T. B., Worsnop, D. R., Malm, W., Wold, C. E., Hao, W. M., and Collett, J. L.: Chemical smoke marker emissions during flaming and smoldering phases of laboratory open burning of wild-land fuels, *Aerosol Sci. Tech.*, 44, i–v, doi:10.1080/02786826.2010.499884, 2010.

Marple, V. A., Rubow, K. L., and Behm, S. M.: A Microorifice Uniform Deposit Impactor (MOUDI): description, calibration, and use, *Aerosol Sci. Tech.*, 14, 434–446, doi:10.1080/02786829108959504, 1991.

May, A. A., Saleh, R., Hennigan, C. J., Donahue, N. M., and Robinson, A. L.: Volatility of organic molecular markers used for source apportionment analysis: measurements and implications for atmospheric lifetime, *Environ. Sci. Technol.*, 46, 12435–12444, doi:10.1021/es302276t, 2012.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- May, A. A., Levin, E. J. T., Hennigan, C. J., Riipinen, I., Lee, T., Collett, J. L., Jimenez, J. L., Kreidenweis, S. M., and Robinson, A. L.: Gas-particle partitioning of primary organic aerosol emissions: 3. Biomass burning, *J. Geophys. Res.-Atmos.*, 118, 11327–11338, doi:10.1002/jgrd.50828, 2013.
- 5 May, A. A., McMeeking, G. R., Lee, T., Taylor, J. W., Craven, J. S., Burling, I., Sullivan, A. P., Akagi, S., Collett, J. L., Flynn, M., Coe, H., Urbanski, S. P., Seinfeld, J. H., Yokelson, R. J., and Kreidenweis, S. M.: Aerosol emissions from prescribed fires in the United States: a synthesis of laboratory and aircraft measurements, *J. Geophys. Res.-Atmos.*, 119, 11826–11849, doi:10.1002/2014JD021848, 2014.
- 10 McMeeking, G. R., Kreidenweis, S. M., Baker, S., Carrico, C. M., Chow, J. C., Collett, J. L., Hao, W. M., Holden, A. S., Kirchstetter, T. W., Malm, W. C., Moosmüller, H., Sullivan, A. P., and Wold, C. E.: Emissions of trace gases and aerosols during the open combustion of biomass in the laboratory, *J. Geophys. Res.*, 114, D19210, doi:10.1029/2009JD011836, 2009.
- 15 Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of composition-dependent collection efficiencies for the aerodyne aerosol mass spectrometer using field data, *Aerosol Sci. Tech.*, 46, 258–271, doi:10.1080/02786826.2011.620041, 2012.
- 20 Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, doi:10.5194/acp-10-4625-2010, 2010.
- 25 Ortega, A. M., Day, D. A., Cubison, M. J., Brune, W. H., Bon, D., de Gouw, J. A., and Jimenez, J. L.: Secondary organic aerosol formation and primary organic aerosol oxidation from biomass-burning smoke in a flow reactor during FLAME-3, *Atmos. Chem. Phys.*, 13, 11551–11571, doi:10.5194/acp-13-11551-2013, 2013.
- 30 Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmos. Chem. Phys.*, 5, 799–825, doi:10.5194/acp-5-799-2005, 2005.
- Robinson, A. L., Grieshop, A. P., Donahue, N. M., and Hunt, S. W.: Updating the conceptual model for fine particle mass emissions from combustion systems, *J. Air Waste Manage. Assoc.*, 60, 1204–1222, doi:10.3155/1047-3289.60.10.1204, 2010.

**Observations and
analysis of organic
aerosol evolution in
some prescribed fire
smoke plumes**

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Schwarz, J. P., Gao, R. S., Fahey, D. W., Thomson, D. S., Watts, L. A., Wilson, J. C., Reeves, J. M., Darbeheshti, M., Baumgardner, D. G., Kok, G. L., Chung, S. H., Schulz, M., Hendricks, J., Lauer, A., Kärcher, B., Slowik, J. G., Rosenlof, K. H., Thompson, T. L., Langford, A. O., Loewenstein, M., and Aikin, K. C.: Single-particle measurements of midlatitude black carbon and light-scattering aerosols from the boundary layer to the lower stratosphere, *J. Geophys. Res.*, 111, D16207, doi:10.1029/2006JD007076, 2006.
- Simoneit, B. R. T., Schauer, J. J., Nolte, C. G., Oros, D. R., Elias, V. O., Fraser, M. P., Rogge, W. F., and Cass, G. R.: Levoglucosan, a tracer for cellulose in biomass burning and atmospheric particles, *Atmos. Environ.*, 33, 173–182, doi:10.1016/S1352-2310(98)00145-9, 1999.
- Stephens, M., Turner, N., and Sandberg, J.: Particle identification by laser-induced incandescence in a solid-state laser cavity, *Appl. Optics*, 42, 3726, doi:10.1364/AO.42.003726, 2003.
- Sueper, D., DeCarlo, P. F., Aiken, A. C., and Jimenez, J. L.: ToF-AMS High Resolution Analysis Software, [online] available at: http://cires.colorado.edu/jjimenez-group/wiki/index.php/ToF-AMS_Analysis_Software (last access: 10 March 2014), 2013.
- Sullivan, A. P., Holden, A. S., Patterson, L. A., McMeeking, G. R., Kreidenweis, S. M., Malm, W. C., Hao, W. M., Wold, C. E., and Collett, J. L.: A method for smoke marker measurements and its potential application for determining the contribution of biomass burning from wildfires and prescribed fires to ambient PM_{2.5} organic carbon, *J. Geophys. Res.*, 113, D22302, doi:10.1029/2008JD010216, 2008.
- Sullivan, A. P., May, A. A., Lee, T., McMeeking, G. R., Kreidenweis, S. M., Akagi, S. K., Yokelson, R. J., Urbanski, S. P., and Collett Jr., J. L.: Airborne characterization of smoke marker ratios from prescribed burning, *Atmos. Chem. Phys.*, 14, 10535–10545, doi:10.5194/acp-14-10535-2014, 2014.
- Urbanski, S. P.: Combustion efficiency and emission factors for wildfire-season fires in mixed conifer forests of the northern Rocky Mountains, US, *Atmos. Chem. Phys.*, 13, 7241–7262, doi:10.5194/acp-13-7241-2013, 2013.
- Urbanski, S. P., Hao, W. M., and Nordgren, B.: The wildland fire emission inventory: western United States emission estimates and an evaluation of uncertainty, *Atmos. Chem. Phys.*, 11, 12973–13000, doi:10.5194/acp-11-12973-2011, 2011.
- Vakkari, V., Kerminen, V.-M., Beukes, J. P., Tiitta, P., van Zyl, P. G., Josipovic, M., Venter, A. D., Jaars, K., Worsnop, D. R., Kulmala, M., and Laakso, L.: Rapid changes in

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

biomass burning aerosols by atmospheric oxidation, *Geophys. Res. Lett.*, 41, 2644–2651, doi:10.1002/2014GL059396, 2014.

Ward, D. E. and Radke, L. F.: Emissions measurements from vegetation fires: a comparative evaluation of methods and results, in: *Fire in the Environment: The Ecological, Atmospheric, and Climatic Importance of Vegetation Fires*, edited by: Crutzen, P. J. and Goldammer, J. G., John Wiley & Sons, Inc., Chichester, England, 53–76, 1993.

Watson, J. G., Chow, J. C., Chen, L.-W. A., Lowenthal, D. H., Fujita, E. M., Kuhns, H. D., Sodeman, D. A., Campbell, D. E., Moosmüller, H., Zhu, D., and Motallebi, N.: Particulate emission factors for mobile fossil fuel and biomass combustion sources., *Sci. Total Environ.*, 409, 2384–2396, doi:10.1016/j.scitotenv.2011.02.041, 2011.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire INventory from NCAR (FINN): a high resolution global model to estimate the emissions from open burning, *Geosci. Model Dev.*, 4, 625–641, doi:10.5194/gmd-4-625-2011, 2011.

Wiedinmyer, C., Quayle, B., Geron, C., Belote, A., McKenzie, D., Zhang, X., O'Neill, S., and Wynne, K. K.: Estimating emissions from fires in North America for air quality modeling, *Atmos. Environ.*, 40, 3419–3432, doi:10.1016/j.atmosenv.2006.02.010, 2006.

Wilson, J. C., Lafleu, B. G., Hilbert, H., Seebaugh, W. R., Fox, J., Gesler, D. W., Brock, C. A., Huebert, B. J., and Mullen, J.: Function and performance of a low turbulence inlet for sampling supermicron particles from aircraft platforms, *Aerosol Sci. Tech.*, 38, 790–802, doi:10.1080/027868290500841, 2004.

Yokelson, R. J., Urbanski, S. P., Atlas, E. L., Toohey, D. W., Alvarado, E. C., Crouse, J. D., Wennberg, P. O., Fisher, M. E., Wold, C. E., Campos, T. L., Adachi, K., Buseck, P. R., and Hao, W. M.: Emissions from forest fires near Mexico City, *Atmos. Chem. Phys.*, 7, 5569–5584, doi:10.5194/acp-7-5569-2007, 2007.

Yokelson, R. J., Crouse, J. D., DeCarlo, P. F., Karl, T., Urbanski, S., Atlas, E., Campos, T., Shinozuka, Y., Kapustin, V., Clarke, A. D., Weinheimer, A., Knapp, D. J., Montzka, D. D., Holloway, J., Weibring, P., Flocke, F., Zheng, W., Toohey, D., Wennberg, P. O., Wiedinmyer, C., Mauldin, L., Fried, A., Richter, D., Walega, J., Jimenez, J. L., Adachi, K., Buseck, P. R.,

Hall, S. R., and Shetter, R.: Emissions from biomass burning in the Yucatan, Atmos. Chem. Phys., 9, 5785–5812, doi:10.5194/acp-9-5785-2009, 2009.

5 Yokelson, R. J., Burling, I. R., Gilman, J. B., Warneke, C., Stockwell, C. E., de Gouw, J., Akagi, S. K., Urbanski, S. P., Veres, P., Roberts, J. M., Kuster, W. C., Reardon, J., Griffith, D. W. T., Johnson, T. J., Hosseini, S., Miller, J. W., Cocker III, D. R., Jung, H., and Weise, D. R.: Coupling field and laboratory measurements to estimate the emission factors of identified and unidentified trace gases for prescribed fires, Atmos. Chem. Phys., 13, 89–116, doi:10.5194/acp-13-89-2013, 2013.

ACPD

15, 1953–1988, 2015

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.




Figure 1. Flight tracks colored by (a) total OA concentration and (b) excess OA concentration. Due to the log-scaling of intercept-averaged concentrations, the minimum value in panel (b) is set to $0.1 \mu\text{g m}^{-3}$. Removing the background OA elucidates distinct plume transport to the southwest.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.




Figure 2. Near-source and downwind data collected during the FJ 9b prescribed fire. **(a)** the ratio of excess OA to CO; **(b)** the ratio of excess rBC to CO; **(c)** emission/export factor for CO; and **(d)** emission/export factor of CO_2 . Near-source data are represented by box-and-whisker plots (boxes: 25th and 75th percentiles; whiskers: 10th and 90th percentiles; horizontal lines: median) while downwind data are represented by markers (closed markers: Lagrangian data; open markers: non-Lagrangian data). Error bars associated with the markers indicate range of estimated time since emission (x direction) and measurement uncertainty (y direction). Error bars in x direction do not account for estimated 30% accuracy of windspeed.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.




Figure 3. As in Fig. 2, but for the Francis Marion prescribed fire.

[Title Page](#)

[Abstract](#) | [Introduction](#)

[Conclusions](#) | [References](#)

[Tables](#) | [Figures](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

[Back](#) | [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.




Figure 4. Evolution of background-corrected AMS mass fractions. **(a)** Δf_{60} for the FJ 9b fire; **(b)** Δf_{44} for the FJ 9b fire; **(c)** Δf_{60} for the Francis Marion fire; **(d)** Δf_{44} for the Francis Marion fire. In all panels, there is a statistically-significant difference between data collected near the source and downwind. Box-and-whisker plots and markers are identical to those in Figs. 2 and 3.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.




Figure 5. Evolution of elemental ratios derived from AMS data. **(a)** O : C for the FJ 9b fire; **(b)** H : C for the FJ 9b fire; **(c)** O : C for the Francis Marion fire; **(d)** H : C for the Francis Marion fire. For both fires, changes in O : C with increasing estimated time since emission are statistically significant. Dashed line is the value of the parameter in the background measurements outside of plume penetrations. Box-and-whisker plots and markers are identical to those in Figs. 2 and 3.

Observations and analysis of organic aerosol evolution in some prescribed fire smoke plumes

A. A. May et al.




Figure 6. Changes in the emission factor of excess OA due to gas-particle partitioning as a function of total observed OA. **(a)** near-source (circles) and downwind (squares) data for the FJ 9b fire; **(b)** near-source (circles) and downwind (squares) data for the Francis Marion fire; **(c)** near-source data for the FJ 22b (upward-facing triangles) and Georgetown (downward-facing triangles) fires; and **(d)** near-source data for the two fires attributed to the Bamberg site (“A”: upward-facing triangles; “B”: downward-facing triangles).