7-2009

SUM OF "N" CONSECUTIVE INTEGERS

Steve Humble

Follow this and additional works at: <https://scholarworks.umt.edu/tme>

Part of the [Mathematics Commons](#)

Let us know how access to this document benefits you.

Recommended Citation

Humble, Steve (2009) "SUM OF "N" CONSECUTIVE INTEGERS," *The Mathematics Enthusiast*. Vol. 6 : No. 3 , Article 14.

DOI: <https://doi.org/10.54870/1551-3440.1167>

Available at: <https://scholarworks.umt.edu/tme/vol6/iss3/14>

This Article is brought to you for free and open access by ScholarWorks at University of Montana. It has been accepted for inclusion in The Mathematics Enthusiast by an authorized editor of ScholarWorks at University of Montana. For more information, please contact scholarworks@mso.umt.edu.

Sum of n Consecutive Numbers

Steve Humble¹

The National Centre for Excellence in the Teaching of Mathematics, UK

Theorem

For all n , it is always possible to find at least one sum of n consecutive numbers with an equivalent sum of $n - 1$ consecutive numbers?

Until recently I did not realise that this wonderful pattern existed.

$$\begin{aligned}1+2&=3 \\4+5+6&=7+8 \\9+10+11+12&=13+14+15 \\16+17+18+19+20&=21+22+23+24 \\&\text{etc}\end{aligned}$$

My first thoughts on reading this connection in O'Shea's[1] book about number curiosities, was why had I not read about it before? It is simply beautiful. O'Shea writes but a few lines on it, and then moves on to his next strange fact. This made me wonder if the pattern would always be true, and here is my proof that it is.

Note that the LHS always starts with a square number. This will always be true, as square numbers occur in the natural number system as follows; 1, 4, 9, 16, 25, 36... with a common difference of 3, 5, 7, 9.....which you can see fits the pattern above. Therefore we can say that each line will always start with a square number. The first few lines in the pattern can be shown to be true, hence it can be proven that the pattern is true for all natural numbers, by considering that the next k th line in the pattern is true.

¹ **Steve Humble** (aka Dr Maths) is a regular contributor to The Montana Mathematics Enthusiast. He works for The National Centre for Excellence in the Teaching of Mathematics in the North East of England (<http://www.ncetm.org.uk>). He believes that the fundamentals of mathematics can be taught via practical experiments. For more information on Dr Maths go to <http://www.ima.org.uk/Education/DrMaths/DrMaths.htm>

Email: drmaths@hotmail.co.uk

Humble

$$k^2 + (k^2 + 1) + (k^2 + 2) + \dots + (k^2 + k) = (k^2 + k + 1) + (k^2 + k + 2) + \dots + (k^2 + k + k)$$

To show that LHS equal RHS, collect the k , k^2 terms on both sides

$$k^3 + k^2 + (1) + (2) + \dots + (k) = k^3 + (k + 1) + (k + 2) + \dots + (k + k)$$

Then collect $(1 + 2 + 3 + \dots + k) = \frac{1}{2}k(k + 1)$ on both sides, giving

$$k^3 + k^2 + \frac{1}{2}k(k + 1) = k^3 + \frac{1}{2}k(k + 1) + (k) + (k) + \dots + (k)$$

$$k^3 + k^2 + \frac{1}{2}k(k + 1) = k^3 + \frac{1}{2}k(k + 1) + k(k)$$

Therefore true for k , hence true for all and proof of the above theorem.

Corollary

In each line in this natural number pattern, we find a triangular, square and cube number sequence.

Angel Proof – “as if at a glance”

Between each pair of square numbers there are $2n$ numbers, n on the LHS and n on the RHS of the above pattern. By adding n to each of the n numbers on the LHS we obtain the RHS.

Reference

[1]The magic numbers of the professor by Owen O’Shea & Underwood Dudley [Page 135]
Published by The Mathematical Association of America 2007