

University of Montana

ScholarWorks at University of Montana

Graduate Student Theses, Dissertations, &
Professional Papers

Graduate School

2015

An Analysis of Socioeconomic Influences on Health Care Seeking Behavior in Humla, Nepal

Hannah Kuehl

University of Montana - Missoula

Follow this and additional works at: <https://scholarworks.umt.edu/etd>



Part of the [Social and Cultural Anthropology Commons](#)

Let us know how access to this document benefits you.

Recommended Citation

Kuehl, Hannah, "An Analysis of Socioeconomic Influences on Health Care Seeking Behavior in Humla, Nepal" (2015). *Graduate Student Theses, Dissertations, & Professional Papers*. 4477.
<https://scholarworks.umt.edu/etd/4477>

This Thesis is brought to you for free and open access by the Graduate School at ScholarWorks at University of Montana. It has been accepted for inclusion in Graduate Student Theses, Dissertations, & Professional Papers by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact scholarworks@mso.umt.edu.

AN ANALYSIS OF SOCIOECONOMIC INFLUENCES ON HEALTH CARE

SEEKING BEHAVIOR IN HUMLA, NEPAL

By

HANNAH ELIZABETH KUEHL

Bachelors of Arts, Point Loma Nazarene University, San Diego, CA, 2009

Thesis

Presented in partial fulfillment of the requirements
for the degree of

Master of Arts
in Cultural Anthropology

The University of Montana
Missoula, MT

May 2015

Approved by:

Sandy Ross, Dean of The Graduate School
Graduate School

Kimber McKay, Chair
Department of Anthropology

Teresa Sobieszczyk
Department of Sociology

Richard Sattler
Department of Anthropology

© COPYRIGHT

by

Hannah Elizabeth Kuehl

2015

All Rights Reserved

Kuehl, Hannah, Masters of Arts, May of 2015, Anthropology

Abstract: An Analysis of Socioeconomic Influences on Health Care Seeking Behavior in Humla, Nepal

Chairperson: Kimber McKay

Abstract: Mobile medical camps (MMCs) are common in Nepal, as well as other parts of the world. MMCs in Nepal provide health care resources that are not available through the government health system, due to Nepal's weak health infrastructure. Although MMCs are problematic for a variety of reasons, they continue to play a role in health care delivery in remote settings the world over. Despite the fact that MMCs are increasingly common in many countries around the world, their patient populations and their health outcomes are poorly understood. This thesis is an analysis of socioeconomic influences within a select group of patients who sought health care from MMCs in Humla district of northwest Nepal. This thesis was written with the goal of better understanding how a person's socioeconomic status (SES) affects their health care seeking behavior via the vehicle of mobile medical camps.

Acknowledgements

I would like to thank Kimber McKay and the Anthropology Department at University of Montana in Missoula for the opportunity of higher education and research in the field of cultural anthropology. In addition, I would like to thank Teresa Sobieszczyk and Richard Sattler for participating in my thesis committee. I would also like to thank Catherine Sanders for her valuable insight and help with the database used for my performed analyses. Matthew Nordhagen also provided help with the data management program, ArcGIS, and for that I am grateful. Randy Skelton taught me the ups and downs of SPSS and helped me better formulate my quantitative analyses.

Table of Contents

	Page
Abstract Page	iii
Acknowledgements	iv
Table of Contents	v
List of Acronyms & Abbreviations	vii
List of Tables	viii
List of Figures	x
Introduction	1
The Rural Region of Humla, Nepal	1
Theoretical Concepts to Consider	5
Impacting Rural and Economically Undeveloped Regions Through this Research	7
Review of Literature: Barriers that Prevent Accessible Health Care	10
Socioeconomic Barriers that Prevent Accessible Health Care	15
Humla’s Weak Health Infrastructure	20
The Rise of Mobile Medical Camps (MMCs)	22
Data Acquisition and Preparation	30
Limitations in Data Acquisition	31
Analysis Methodology	32
Descriptive Statistical Analysis	34
Inferential Statistical Analysis	35
Spatial Examination of Health Care Seeking Routes	36
Results of Statistical Analyses and Spatial Construction of Routes	37

Discussion	49
Suggestions for Further Research	51
Implications for MMCs in Health Care	53
References	55

List of Acronyms & Abbreviations

ArcGIS	Geographic Information System
DHS	Demographic Health Survey (Wealth Index)
INGO	International Nongovernmental Organization
MMC	Mobile Medical Camp
NGO	Nongovernmental Organization
SES	Socioeconomic Status
SPSS	Statistical Package for the Social Sciences
UNFCO	United Nations Field Coordination Office

List of Tables

Table	Page
Table 1: Patient Chief Medical Complaint	37
Table 2: Patient Socioeconomic Status	37
Table 3a: Patient Gender and Age in Years (Age 1-50)	38
Table 3b: Patient Gender and Age in Years (Age 51-90)	38
Table 4: Patient Socioeconomic Status and MMC Attended	38
Table 5: MMC (Chauganphaya) Attendance Based Upon Village of Patients' Residence	39
Table 6: MMC (Dandaphaya) Attendance Based Upon Village of Patients' Residence	39
Table 7: MMC (Kermi) Attendance Based Upon Village of Patients' Residence	39
Table 8: MMC (Muchu) Attendance Based Upon Village of Patients' Residence	40
Table 9: MMC (Syada) Attendance Based Upon Village of Patients' Residence	40
Table 10: MMC (Yalbang) Attendance Based Upon Village of Patients' Residence	40
Table 11: Patient Childcare Decision Making	40
Table 12: Patient Childcare Provider	41
Table 13: Patient Work Replacement	41
Table 14: Amount of Time Traveled to MMC (in minutes), excluding outliers	41
Table 15: Test for Normal Distribution of Dependent Variables	44

Table 16: Homogeneity of Variances for SES	45
Table 17: Descriptive Statistics of one-way ANOVA for SES	45
Table 18: Results for One-way ANOVA Analysis for SES and Dependent Variables Tested	46
Table 19: Chi-Square Test between SES and Variables	48

List of Figures and Illustrations

Figure	Page
Figure 1: A map of Nepal with Humla district in red	2
Figure 2: Test for Outliers within Number of Adults in Household	42
Figure 3: Test for Outliers within Amount of Time Traveled to MMC	43
Figure 4: Test for Outliers within When Previous Treatment was Received (in months)	43
Figure 5: Walking routes from villages to MMCs	49

Introduction

Humla district is a rural and poor region of Nepal, stranded in the Himalaya mountains (Justice 1986, 5). It lacks infrastructure in many areas, specifically health care (Adara 2014). NGOs are common in this region of Nepal, providing health care, education, and sanitation technologies to the residents (Sanders and McKay 2013, 109). One NGO conducted a qualitative improvement (QI) study in this region, using a Mobile Medical Camp (MMC) to deliver health care. The study focused on gathering data to gain information on how better it can improve its services. As part of my thesis, I used this de-identified data to conduct analyses to identify socioeconomic barriers on health care seeking behavior.

The Rural Region of Humla, Nepal

Nepal is among the poorest countries in the world (Citrin 2011, 16). The 2011 Human Development Index (HDI) indicates that as much as 40% of Nepal's population is below the poverty line. Nepal's poverty rate was ranked 157th out of 187 countries in the world in 2011 (Daniel 2012, 6-7).

Nepal is divided into three zones: the northern Himalayas (the mountains), the central hills (the hills), and the flat southern plains (the Terai) (Daniel 2012, 7). The district of Humla lies between 81 and 82 degrees longitude and 30 degrees N in latitude (McKay 2002, 241) in the far northwestern corner of Nepal in the central Himalayas (Justice 1986, 5). It is bordered by China, which was formerly Tibet (Grocke 2013, 1). Figure 1 shows a map of Nepal. Humla district is shown in red.




Figure 1: A map of Nepal with Humla district in red (Wikipedia 2014)

Within Nepal, Humla district is at a disadvantage, being isolated from the rest of the country in the high mountains (Adara 2014). “A peripheral area in a peripheral country, Humla is a vast and road-less mountain region that consistently ranks at or near the bottom of Nepal’s 75 districts on many socio-demographic and health parameters” (Citrin 2011, 20). In 2011 it had the lowest literacy rate in the country; 47.8% compared to a national average of 65.9% (Central Bureau of Statistics 2012, 4, 226-260).

Different crops are grown seasonally: rice, maize, and potatoes from April to June, wheat from October to November, and apples from December to January. The summer and growing seasons are short and affected by avalanches, floods, landslides, fire, storms, droughts, glacial lake outbreaks, and rocky terrain. The region’s high elevation (1,524 - 7,337 meters) and geographic conditions lead to a short growing

season and high food insecurity (UNFCO 2013, 2, 10-11). The winter is long and harsh with temperatures ranging from 25 to -28 degrees Celsius (UNFCO 2013, 11).

There are no roads in the entire district or in connection “to the rest of the country” yet, although the Hilsa-Simkot road is currently under construction. This road will eventually connect Humla district to Tibet (UNFCO 2013, 1-2). Simikot, the district headquarters, is only accessible by foot or airplane. Because planes are unreliable due to weather conditions and unaffordable for most residents, most areas in Humla are only accessible by foot and “can be reached only by walking for hours or even days along steep, winding trails” (Justice 1986, 5).

The Maoist conflict, which is now largely over and lasted from 1996 to 2006 (Basnett 2009, 4), has had a large impact on the difficulty of travel in remote places like Humla (cf. Citrin 2012, 74). The main Humla-Karnali River, as well as 50 other smaller rivers, divide Humla district. The rivers “...are uncrossable when swollen by monsoon rains and melting snows, [and] they are during much of the year” (Justice 1986, 5). There used to be 27 suspension bridges connecting different parts of Humla together, however, the Maoist conflict destroyed many bridges during the war. In 2013, many bridges were still not functional (UNFCO 2013, 1-2).

Due to the geography and climate, the elevation, and the loss of bridges, most villages are cut off from the outside world for up to four months out of the year (UNFCO 2013, 11). Development of infrastructure, including transportation, communication, and health care, has been hindered due to the isolation of rural areas from big cities and government. Geographic distance, as well as the mountainous terrain, provides difficult barriers to overcome. This leads to the division and isolation of rural areas (Justice 1986,

5). Infrastructure is poor, education is limited, and paid employment is low due to the prevalence of a non-cash economy (Adara 2014).

As with many other economically developing countries, Nepal endures health challenges (Dupas 2011, 2). However, Humla seems to be at a greater disadvantage than even the rest of Nepal. Preventable and infectious illnesses are amongst the highest in the country. Over 60% of the district's population suffers from stunted growth and 15% are severely underweight. Fifty-three percent of children below five years of age suffer from anemia (UNFCO 2013, 9). Food shortages are common; only 10% of the district has enough food for an entire year (UNFCO 2013, 11).

The main health concerns are severe malnutrition, critical respiratory infections, childbirth, and complications due to heavy worm loads and Hepatitis B, which is endemic in this population. Respiratory diseases are common due to smoking and indoor air pollution caused from fire-burning stoves. Lack of sanitation and hygiene technologies lead to food and water-borne diseases. Steep mountain paths produce injuries from falls and landslides. The local subsistence system is based on the practice of raising livestock, agriculture and long distance pedestrian trade. The work is taxing and only exacerbates the burden of disease (Sanders and McKay 2013, 110; UNFCO 2013, 8).

For most of these diseases and ailments there are appropriate treatments, yet still these diseases continue to cause morbidity and mortality. Worldwide, 6-70% of child deaths are due to deficiency in seeking health care (O'Donnell 2007, 2821). People in economically undeveloped countries are strongly affected by infectious and parasitic diseases that render receiving health care of great importance in order for children and

adults to lead as long and healthy lives as people in more economically developed countries (Dupas 2011, 2).

Theoretical Concepts to Consider

“The ‘quest for therapy’ all over the world is an important research issue since it reveals essential elements of people’s social behavior and provides insights into their perceived needs for different kinds of health services” (Kroeger 1983, 147). This research will draw upon the theoretical framework of critical medical anthropology and focus specifically on the political economy of health. It will draw upon a conceptual model within the greater social sciences: the health behavioral model. It will combine the political, economic, social and cultural determinants of health care seeking behavior to provide a rounded theoretical investigation.

Critical medical anthropology is founded upon the idea that “change programs should not be attempted unless one also studies the social production of illness and poverty within the larger dynamics of class interactions, colonialism, or world economic systems” (McElroy 1996, 7). Marxist and dependency theory both influenced critical medical anthropology in its analysis of “...the impact of global economic systems, particularly capitalism, on local and national health” (McElroy 1996, 7). As Baer has suggested, access to health care is influenced by cultural and historical conditions, as well as hegemonic systems of power. The idea of accessible health care depends upon “how and in what form... this power [is] delegated, and how this power is expressed” (Alexandrakis 2001, 75). In order to study health care seeking behavior, it is crucial that first the “political, economic, social and environmental conditions [are identified] within a particular society or group” (Alexandrakis 2001, 76).

Across the field of social science, social psychology's health behavior model is one of the oldest and best known (Hausmann-Muela et al. 2003, 9). This model suggests that health care behavior is a combination of 4 factors: perceived susceptibility, perceived severity, perceived benefits and perceived barriers (Coreil 2008, 76-77). Hyder and Morrow suggest that the strongest predictive factor on health care behavior is perceived barriers (2005, 51). Focusing on perceived barriers as the strongest predictive factor on health care behavior makes the health behavior model relevant in measuring the costs and benefits of accessing health care services (Coreil 2008, 76-77).

Although the health behavior model is most often used within the realm of health promotion, it can be applicable towards understanding individual health seeking behavior patterns. According to Mackian, this model's downfall lies in assuming the decision-maker is a purely rational being (2003, 8). As Ward, Mertens, and Thomas has suggested, it is not rational to assume "that recognition of symptoms will necessarily or automatically result in health seeking behavior" (Bork-Huffer 2012, 25).

This thesis will use analyses to show patterns of association, however, it will not underestimate an individuals' capacity for agency and resistance (De Los Angeles Nunez Carrasco 2008, 40). It is important, especially as an anthropologist, to remember, "health and illness are defined, labeled, evaluated, and acted upon in the context of culture" (Hyder, Morrow 2005, 41). The health behavior model also "neglects determinants... like previous experiences, advantages of mal-adaptive behavior, behavioral intention, perceived control etc." (Hausmann-Muela et al. 2003, 10). Although this thesis will be based upon quantitative data and analyses, qualitative search for meaning and reasoning will be pursued to explain possible determinants.

Impacting Rural and Economically Undeveloped Regions Through this Research

The health infrastructure is weak in Nepal. Mobile medical camps (MMCs) are common there, as well as other parts of the world. They function to provide health care resources that are not available through the government system of sub health posts, health posts and district and regional hospitals. Although MMCs are problematic for a variety of reasons, they continue to play a role in health care delivery in remote settings the world over.

In order for Humli residents to become healthier, NGOs working in the region believe health care seeking behavior must evolve. In order for health care seeking behavior to change, we need to better understand current health seeking behavior patterns. This project is an analysis of one such attempt by an NGO to gather health and social information about patients, in order to create a lasting and long-term health partnership with this community and individuals within it.

MMCs and their outcomes are poorly understood. In order to broaden our understanding of how an MMC might be perceived by a local person, an NGO working in remote Humla decided to collect not just clinical data, but also social, economic, and attitudinal data from patients. Because these data address health care seeking behavior, this is primarily the literature that I have engaged with. This thesis is a study and analysis of data collected by the NGO staff on patient socioeconomics and logistics of travel.

The first set of analyses presented here explores whether socioeconomic status (SES) has an effect on health care seeking behavior. This section of the project addresses the household level trends and barriers against seeking health care. Results show how key socioeconomic factors affect health care decision making within a group of individuals

seeking health care from local government and private clinics as well as MMCs in the Humla district of Nepal.

The second set of analyses presented in this thesis represents an attempt to better understand the logistics of travel needed to obtain health care in a remote and mountainous region, such as Humla. Although health care seeking behavior is heavily “influenced by the distance [the patient has] to travel... few studies have tried to estimate these distance effects” (Muller et al. 1998, 878). Jolly and King were the first to quantitatively demonstrate the effect that travel distance has on the utilization of health care services (Kroeger 1983, 151-2). This thesis will explore travel routes, distances, and elevation gains and losses required for villagers to access health care. The primary intent behind these analyses is to better understand the role of distance decay (the decrease in utilization with distance) in local people’s health seeking behavior (Mueller et al. 1998, 878).

Results from these analyses may add towards a greater understanding of health care seeking behavior. Ultimately, results from this thesis can be of aid to the major issue “how can the benefits available to the few be expanded to meet the needs of the many?” (Balabanova et al. 2006, 2). Results may help NGOs be able to provide more adequate services in Northwestern Nepal. Specifically, results may propose better locations for MMCs in Humla district. These revised locations will make health care more easily travelable and accessible for patients.

Results may have broader implications for similar geographic regions. Findings related to socioeconomic factors and travel logistics of seeking health care can be used for geographic regions that have similar climates and health care resource limitations. An

important part of providing successful local health care is the study of how people use health care facilities (Kroeger 1983, 147). The results of these analyses will be helpful to villagers located in rural and high elevation areas and to the NGOs working within similar regions all around the world. This thesis benefits the villagers within Humla district, the broader community, and the larger anthropological community.

I have already discussed the geographic vulnerabilities of Humla, Nepal. Next, I review literature based upon the barriers that prevent people from seeking health care. It will focus specifically on socioeconomic barriers. Then I discuss the weak health infrastructure in Humla. Finally, I will talk about Mobile Medical Camps (MMCs) and address their pros and cons on the community.

After the literature review, in order to better understand the relationship that socioeconomics has with health care seeking behavior, I will report results from data that were collected from six MMCs in northern Humla in 2013. To show hardship of travel and inaccessibility of health care access, I will show a map of 26 commonly used routes in Humla district. I will conclude with a discussion of my results.

It is important to define *health care seeking behavior*, as this term can cause confusion. It is "...conceptualized as a 'sequence of remedial actions' taken to rectify 'perceived ill-health' (Ahmed et al. 2000)" (Mackian 2003, 7). *Health care seeking behavior* can include both formal and informal methods of seeking health care. This is different from *health seeking behavior*, which is more heavily adopted in psychology and looks at the general idea of what it means to be healthy (Mackian 2003, 7). *Health seeking behavior* focuses on health-promoting behaviors as opposed to actions one takes

when one is sick. Due to the database I worked with, the focus of this thesis is *health care seeking behavior*. This focus is most appropriate for the data being analyzed.

Review of Literature: Barriers that Prevent Accessible Health Care

Convincing amounts of literature demonstrate that the issue of health care accessibility continues to be a prominent problem in economically developing countries (O'Donnell 2007, 2821, 2830). However, accessible health care has come to have different meanings for different people. There are many definitions that relate to the term *access to health care*. Sometimes access to health care is only defined in its reference to geography. Penchansky and Thomas refer to it as a combination of four individual parts: affordability, availability, acceptability, and accessibility. While some people understand it to mean "... the opportunity to use health care; others draw no distinction between access and use" (O'Donnell 2007, 2821). O'Donnell "proposes that the central concern is whether individuals that can potentially benefit from effective health care do in fact receive it" (2007, 2821).

Barriers to health care are categorized as either supply-side or demand-side. Ensor and Cooper believe that demand-side barriers are determinants of health care use that are not dependent on price or service delivery. They would consider cultural and social barriers, education, opportunity cost, and distance all to be considered demand-side barriers (Ensor, Cooper 2004, iii). Barriers to accessible health care are often not clear because they "...may interact and influence each other" (Jacobs et al. 2012, 290). A barrier might act as both a supply and demand-side barrier in different ways. This thesis will first address demand-side barriers that prevent people from accessing health care. It will then look at MMCs as a potential solution to supply-side barriers.

A person's resources, or lack thereof, can act as a barrier to accessing health care. There are many forms of resources that a person might need in order to seek out health care; these include material resources, human resources, and social resources. Each of these factors has an effect on one's ability to access health care in different ways. Material resources can include physical, natural, and financial capital. Physical resources include transportation. Natural resources include ownership of livestock or land, which lends to being wealthier. Financial resources include cash or credit. Human resources include one's level of education and ability to obtain information. Social resources include one's social support network that can include family, friends, and neighbors who are willing to make sacrifices to help a person whom is seeking health care (Bakeera et al. 2009, 6-10). Due to the data available, this thesis will focus on material resources that help to build a person's SES; primarily including natural and financial capital.

It is hard to measure the reasons behind health care seeking behavior. It is much easier to measure the quantifiable effects (e.g. who is accessing health care and what health care is being offered). Oliver and Mossialos argue that scientists oftentimes put more effort into researching the *availability* of health care instead of the *utilization* of health care. The utilization of health care can be unquantifiable and messy. Tipping and Segall understand utilization to be more easily measured and "...a frequent proxy for access" (Balabanova et al. 2006, 2).

Information, reliability and reputability of the information provider, ability to process information, education, and time to adapt to new information or new technologies are all barriers that must be overcome in seeking health care (Dupas 2011, 9-15, 29). Gender discrimination, socioeconomics, and geographical distance are also critical

factors in one's ability to access health care (Deogaonkar 2004, 1). The weaknesses involved in supplying health care include producing and providing inadequate quality of health care and insufficient resources. The supply-side of health care also errors in inappropriately allocating resources across levels of care, regions, and programs (Dupas 2011, 9-15, 29; O'Donnell 2007, 2826).

The dissemination of information can be achieved through numerous forms, such as social networking (e.g. family, relatives, acquaintances, friends), media (e.g. TV, radio, internet, newspaper), information campaigns (e.g. community information, wall posters), and health care facilities (e.g. doctors, information campaigns from hospitals) (Bork-Huffer, 2012, 181). In a study done by Adelman, Essam, and Leonard in 2009, there was evidence to show that households gained and evaluated different health care facilities by watching and learning from their neighbors' experiences, or by using social learning skills (Dupas 2011, 15-16).

Unfortunately, the spread of information does not happen in areas where the coverage of treatment is low because there "...is less opportunity to learn of its benefit" (O'Donnell 2007, 2826). A study to research limitations in health care access within the Saptari district of Nepal showed that most residents living in this community lacked information on medicines and health care services and incentives (Daniel et al. 2012, 3). Most importantly, information was not being disseminated in effective ways.

Reliability and reputability of the organization providing information is critical. The community must trust the provider in order to want to receive health care and seek out health care treatment from the respective source. Trust can be gained or lost by observing the effectiveness of services. If the services provided are successful and high

quality, trust will be established (O'Donnell 2007, 2826). Alderman and the World Bank have done studies that prove demand for health care will go down when quality diminishes. A study in a rural region of India, done by Gwatkin with the World Bank, showed a correlation between low utilization of a free, public health care facility and low quality of health care being provided (O'Donnell 2007, 2081-2).

Members of one's social network highly influence a sick person's ability to access health care. Research participants in a study in the Saptari district of Nepal emphasized the importance of social capital in accessing health care. The utilization of social networks through friends, family and relationships, were shown to be critical in gathering information and support relating to health care, services, and financial help (Daniel et al. 2012, 3). One way support can be provided is by taking over the sick person's chores, work, or housework. Spouses mainly take over this responsibility, however, other family members, such as parents and children, also help in this capacity. Less often, one's neighbors and friends take part in helping to alleviate the burden of work (Bork-Huffer 2012, 242).

Gender inequality is a major factor in reducing access to health care for females in many economically developing countries, including Nepal (Sreeramareddy, Harsha Kumar, Sathian 2013, 2). The Seva Canada Society and Seva Foundation found that "women of all ages (including children) are more frequently exposed to causative factors, such as infectious diseases and malnutrition..." yet they oftentimes have less ability to get transportation to and from health care facilities, have access to less money, and are more geographically isolated (Unite for Sight, 2000-2015(b), 1).

In economically developing countries, the root cause of gender disparities in access to health care can be “traced to the low social status of women in much of the developing world” (Unite for Sight, 2000-2015(b), 2). A study pertaining to eye care produced by Geneau, Massac, Courtight, and Lewallen in Tanzania found “that men were more likely than women to adopt a ‘sick role’. In contrast, women were more likely to continue their routines through ‘adversity’ instead of emphasizing the need for [eye] surgery” (Unite for Sight, 2000-2015(b), 2). In Humla district, gender inequality is worsened by a high incidence of gender-based violence, triggered in part due to a wide availability of Chinese alcohol (UNFCO 2013, 5).

Although females are at a disadvantage in their ability to access health care, studies show that females attach a greater priority to child and maternal health care than males (O’Donnell 2007, 2829). When compared to males, females spend more of their income on health care for their children (Dupas 2011, 13). Although females are at a higher disadvantage in accessing health care, they might seek out resources more than men due to their values and “might respond different[ly] to information on the health returns of various behaviors” (Dupas 2011, 13). Increasing the social conditions of women, including changing gender roles and attitudes, is important in the fight for providing global, accessible health care.

Although it is crucial that maternal, reproductive, and child health care be made more accessible for females, the implementation of health care in this arena remains challenging “within societies that restrict the public lives of women” (O’Donnell 2007, 2826). The cost of work replacement (both paid and unpaid) is a barrier in seeking health care. A study was done by Banerjee, Duflo, Glennerster, and Kothari in 2010 that studied

the effects of health care seeking behavior when the cost of treatment was incentivized by off-setting the cost of losing a day's wage when receiving treatment. The study focused on reliable immunization camps in Udaipur, India. Parents who brought their children to one of the immunization camps received 1 kg of lentils per immunization administered. This was equal to three-quarters day wage for the parent. The incentivized immunizations grew immunization rates from 17% to 38% (Dupas 2011, 25).

Seeking health care for a mother is a burden to herself and also for her children. When seeking health care for herself or one of her children, mothers often need to find childcare for the rest of the children. This requires support from a social network. If the health care facility is far away, the social request for childcare services is much greater (MacKian 2003, 12-13). Childcare is a critical consideration for mothers and women who take care of children (Ward 2009, 1591). Even if women require health-care services, the competing demands of taking care of children can make it impossible to do so (Creel et al. 1996, 3).

Socioeconomic Barriers that Prevent Accessible Health Care

In this project, I felt compelled to generate a better understanding of the impact of socio-economic status (SES) on health care seeking behavior. I analyzed this in a fairly fine-grained way, by looking at behavioral patterns of patients seen at one of the six MMCs I analyzed. As Subedi argues, health issues in Humla , “among men [and] women, cannot be separated from the larger social, cultural, economic, and political forces that shape and constrain human life” (2010, 22). A patient's SES proves to have a large effect on both health care seeking behavior and the health of a population. “Poverty, which is a result of social and economic inequality in a society, is detrimental to the health of

population. The outcome indicators of health (mortality, morbidity and life expectancy) are all directly influenced by the standards of living of a given population” (Deogaonkar 2004, 1).

It is important that we take time to correctly understand the usage of SES as it will be used in this paper. “SES is a difficult concept to define and to measure” (van Bodegom et al. 2009, 68). It can be composed of different indicators, depending upon whether the setting is in economically developed or developing countries. For this paper, SES will be interpreted

as a hierarchical position that determines someone’s access to human or material resources. SES largely influences health and survival in both developing and developed countries. In developed countries SES is mostly measured as a combination of education, income and occupation. In developing countries however, especially in rural areas, most people have little education and income tends to be fluctuant depending both on the season and on the year and is not easily expressed in numbers. Also, the SES in developing countries is largely dependent upon a relative, social position in the community, clan or extended family and this is not always reflected in property. [van Bodegom et al. 2009, 68]

Wilkinson suggests that it is not necessarily the deprivation of income, but instead the distribution of income that matters. Multiple studies done across the globe have revealed a strong association between excess mortality and income inequality. A study done by Kennedy, Kawachi, and Prothrow-Stith, in the United States, showed that

income inequality directly affected the total mortality of the population. This study measured income inequality using the 'Robin Hood Index'. The 'Robin Hood Index' is a measurement of the part of income that would need to be redistributed from the rich to the poor in order to achieve economic equality. By this measure, a 1% rise would lead to 21.7 excess deaths for every 100,000 people (Deogaonkar 2004, 1). It is well known and documented that health disparities exist within Nepal due to inequality in socioeconomics (Sreeramareddy, Harsha Kumar, Sathian 2013, 2).

Demery, in 2000, and Makinen et al., also in 2000, found evidence "...that access to health services and the distribution of public subsidies favor richer, urban dwellers over generally poorer, rural inhabitants" (Ensor, Cooper 2004, 1). Although the poor tend to be the least healthy and need health care the most, they are usually the lowest in utilization of effective health care services. Income disparities, which are greatest in economically developing countries, lead to disparities in access to health care between rich and poor. The poorest 20% of people in Nepal receive less than 10% of the public health subsidy (O'Donnell 2007, 2822- 4, 2830). Research done in the Saptari district of Nepal concluded that socioeconomic factors substantially contributed to limitations in access to health care and poorer health status of Dalit communities (Daniel et al. 2012, 3).

A person's SES affects their level of awareness in relation to health seeking behavior. Lack of awareness is a direct barrier to seeking health care for oneself or one's children. India provides free immunizations to all children, yet still 2 out of 5 children are not fully immunized. Pande and Yazbeck found that although the country provided free immunizations, most female parents did not know about the benefits of immunization. A

large percentage of female parents who did understand the benefits of immunization did not know where to go to receive vaccinations (O'Donnell 2007, 2826).

The ability to recognize illness is essential to seeking health care (Dupas 2011, 9-15, 29). "...Cultural and educational factors may obscure the recognition of illness and the potential benefits from health care..." (O'Donnell 2007, 2821). There is a direct correlation between knowledge of illness and health care utilization. In order to demand health care, there must be an ability to recognize illness and to recognize the benefits of treatment. In a population where poor health is the norm, it is harder to recognize good health as a viable option. Illness is more often reported in people who are wealthy than in people who are poor. "The better off are more likely to seek care for a child when sick, to take anti-malarials and antibiotics for pneumonia, and to receive inpatient care... The unfortunate outcome [for the poor] can be the continued toleration of illness and disease" (O'Donnell 2007, 2826).

Even when a person has full awareness of the availability of health care, the cost of seeking health care treatment might be too great. Cost is oftentimes the most pressing barrier to seeking health care (Unite for Sight 2000-2015(a), 1). Thus, even if health care is recognized as needed, utilization might be suppressed because of economic constraints (O'Donnell 2007, 2821). Gertler and Gaag saw successful health care utilization take place within a poor population in Africa when certain factors were met: the ability to pay with credit, reductions in travel time, and improvements in quality (O'Donnell 2007, 2825). "Individuals must be willing to use effective preventive and treatment interventions and they must have the purchasing power to realize this desire" (O'Donnell 2007, 2827).

There is a positive relationship between one's economic standing and the utilization of health care. Multiple studies have shown a correlation between prenatal care and medically supervised deliveries with a rise in income. Steele, Diamond, Amin and Gage, Sommerfelt, and Piani have shown a positive association between income and child immunization through the use of multivariate analyses (O'Donnell 2007, 2825). "Monetary costs of care [including the non-price costs of long distance travel] ensure that income is an important determinant of health care utilization..." (O'Donnell 2007, 2825)

Cultural beliefs about allopathic medicine and the causes of disease also factor in to one's willingness to seek health care from western clinics and doctors. Fear of procedures is daunting, especially when associated with visiting doctors or by short-term medical interventions (Dupas 2011, 9-15). Social and cultural norms are oftentimes reflected from attitudes that are deeply rooted in family traditions. These can produce a low demand for modern health care. Social norms within groups are constantly evolving. There is a correlation between a rise in education and a decline in seeking traditional health care. Again, the socioeconomic environment influences social norms (O'Donnell 2007, 2826).

Lack of geographical access is a major barrier to receiving effective health care. Distance has shown to have a greater effect on health care utilization in the poor than for wealthy individuals. This makes sense due to poorer people living in areas with worse road conditions, less available transportation, and greater social strains associated with seeking health care. "In rural areas, the distances to health care facilities and the poor condition of roads mean that time, effort, and cost required to arrive at the point of

delivery can be substantial. The evidence confirms the expected negative impact of health care utilization” (O’Donnell 2007, 2826).

A cataract study conducted by Chang, Condon, Baker, Bloem, Savage, and Sommer in rural Africa showed that distance to receiving health care was the most important barrier in 1/3 of all patients (Unite for Sight 2000-2015(a), 1). A study in Ghana showed that when the travel distance to a health care facility was cut in half, the utilization rate almost doubled (O’Donnell 2007, 2826). “Lowering the barrier of distance requires either taking people to services or services to people... road building is expensive and is not under the control of health sector policymakers. More feasible are schemes that lower the price of travel for health care or provide credit to cover travel expenses” (O’Donnell 2007, 2830).

People living in urban areas generally have closer and better geographical access to health care. Cities have the biggest concentration of health care provisions (O’Donnell 2007, 2083- 4). People in rural areas oftentimes have to travel far distances to receive health care and also lack transportation services (Unite for Sight 2000-2015(a), 1). This puts the rural region of Humla, Nepal at a greater disadvantage in being able to access health care than people living in urban settings such as Kathmandu, Nepal.

Humla’s Weak Health Infrastructure

Humla’s population of 50,858 people is supported by only two hospitals and 26 health posts (of which 16 are mostly non-functioning) (UNFCO 2013, 8). The health posts and hospitals frequently lack the appropriate equipment and trained health care personnel (UNFCO 2013, 8). Health posts were weakened during the Maoist conflict (1996-2006) through the ransacking of valuable materials (Citrin 2012, 74). Medical

posts usually consist of untrained staff, expected to treat patients with limited medical education and a small amount of medicines.

All villagers in Humla district live in an extremely rural setting. Health posts are usually not located or in working condition near most of these villages. Furthermore, patients often find the medical posts to be temporarily or permanently closed, after having traveled for an extensive amount of time (Citrin 2011, 34; Justice 1986; Sanders and McKay 2013, 112).

Humla's isolation from the rest of Nepal has attributed to the unavailability of health care (Citrin 2011, 20). The mountainous topography of Humla has limited the infrastructure of health care facilities (Sanders and McKay 2013, 109). In Nepal, there is a major geographical advantage in seeking health care for people who live in or around the Kathmandu Valley. The valley has more plentiful (though still insufficient) doctors and health care resources. To reach Kathmandu from Humla, a walking trek and plane flight is required. Most people who live in Humla cannot afford the transportation costs of a plane flight (UNFCO 2013, 1).

Beyond distance, Humla has the highest elevation of any district in Nepal. Most villages range in elevation between 3,000 to 5,000 meters above sea level (Digital Himalaya's Nepal Maps Collection 2014). Extreme gains and losses in elevation add to an already long trek to seek health care. Long distance and large amounts of elevation gain and loss would make this journey difficult for a sick person. Topography, inadequate resources and shortage of trained medical personnel are all major factors in the inaccessibility of health care within Humla district (Sanders and McKay 2013, 109).

Because Humla district is dependent upon agriculture for most of their food security, seeking health care is highly contingent based upon the growing season. Leatherman's assumption is "if it is harvest or plowing season, time constraints on health seeking are an even greater issue, often delaying treatment processes beyond the critical period" (Sanders and McKay 2013, 112-3). Compensation for a day's loss of work might not be equivalent to the amount of impact that loss of fieldwork will produce. When an individual's health competes with the entire household's economic outcome when faced with the decision to work in the field or seek health care treatment, health concerns are usually overlooked for the good of the family (Sanders and McKay 2013, 116).

The Rise of Mobile Medical Camps (MMCs)

Nepal was not colonized like many of its neighboring South Asian countries, and it remained closed to outsiders and foreign health systems until the 1950s. Western scientific medicine (called allopathic medicine in Asia) was not used in Nepal until a revolution took place that restored the Nepalese monarchy and opened up borders through the creation of an international airport in Kathmandu. Christian missionaries first took advantage of the country's opening to provide health assistance. By 1951, government hospitals were available in Kathmandu (Justice 1986, 8). Before the 1950s, health care was sought through family knowledge and the use of traditional healers. After the opening of the borders, foreign drug companies started utilizing Nepal as an open market for western drugs (Justice 1986, 8).

Although the Nepalese government has made progress in providing medical care for its citizens, most health care advancements had been made in Kathmandu and other cities, as opposed to rural areas where most of Nepal's population still reside (Justice

1986, 9). Furthermore, as determined by the World Bank, scarce resources and inefficiency has neglected to decrease major underlying causes of disease: "...poor nutrition, polluted water, and lack of education in hygiene and sanitation" (Justice 1986, 9).¹

Nongovernment organizations (NGOs) began taking over the government's role in providing health care for the poor and marginalized in the 1950s. The United States, India, the Soviet Union, China and the World Health Organization (WHO) were among the first external health care donors to Nepal (Justice 1986, 8). NGOs today, continue to provide health care services to both rural and poor communities in Nepal. "NGOs provide essential services that would otherwise leave masses of people without basic tools of survival (e.g., sewage disposal, nutritional supplementation, and medicine)" (Sanders and McKay 2013, 109).

Research addressing the politics of health care, health care development, and health care provided by NGOs, has recently began to address their negative aspects (Citrin 2012, 15). The role of short-term, traveling, health care-providing settings, including MMCs, has been included in this critique. I will discuss a critical evaluation of both donor agencies and NGOs within Humla district and MMCs, in particular, because data for this project was obtained through this method of health care.

NGOs in Humla district have come under attack for many legitimate reasons. Beyond the suspicion in motivation for funding, overlapping aid and under met needs, time-related issues, and questions regarding potential benefits to providers instead of

¹ An evaluation of the health care system in Nepal is critical in understanding why NGOs are prevalent in the region. A detailed critique and history of development of the health care system in Nepal is given by Judith Justice in "Policies, Plans, & People: Foreign Aid and Health Development".

people in need, there are theoretical reasons of critique. The ‘medicalization’ of hunger is a major issue that needs addressing.

Funding for NGOs can come from politically motivated organizations. These organizations might not address the most pressing concerns for the community being served, but instead, might try to please the organization that provides the financial backing (UNFCO 2013, 13, Citrin 2012, 18). Donor agencies might have ties associated with funding that can prove more harmful than helpful to the recipient. For example, “USAID is the largest bilateral donor of health aid to developing countries, including Nepal. Its policy guidelines originate in the U.S. Congress, the U.S. Department of State, and the Policy Division of USAID. The agency is therefore accountable to all three levels” (Justice 1986, 26). Within an agency as large and diversified in its interests as USAID, it’s hard to tell whom the agency is truly trying to benefit.

Many times, villagers needs and expectations are completely forgotten, and instead plans are implemented based upon what the donor agency thinks is appropriate (Justice 1986, 33). Therefore instead of being fully accountable to the recipient’s needs, the donor organization is first and foremost accountable to its origins, then to the recipient. Furthermore, to further complicate recipient needs, donor agencies such as USAID hire advisors from private companies, which could have different objectives than even the donor agency (Justice 1986, 28).

Currently, there are more than a hundred NGOs in the district of Humla, Nepal. Better services could be provided if work was consolidated within the organizations instead of duplicated between them. Aid is overlapping in many areas (Justice 1986, 9) and not meeting needs in many others. Even with the vast amount of NGOs available in

Humla district, people still fail to have their most basic health care needs met (Sanders and McKay 2013, 109-110).

Furthermore, competition among donors can create tension for the recipient who is caught in the middle of a funding feud. These battles could be avoided if donor agencies assisted programs created by the Nepalese government, instead of creating their own (Justice 1986, 33). "...Too many donors and too little coordination" (Justice 1986, 33) can lead to massive amounts of confusion and lack of help for the recipient.

Many donor-led projects have to include a time-line and objective to be completed within just a short period of time. Oftentimes the stated objectives are too large or unreasonable to complete. However, to continue receiving aid, material results must be produced. This can lead to falsifying results, or lead to the creation of hard outcomes, even if the falsified outcomes are not what are most beneficial for the recipients (Justice 1986, 27-32).

Funding for projects with short time-lines and hard outcomes lead to a continuous repetition of unfinished and unnecessary projects. Many NGOs leave after completing a faulty and short-term project, even when infrastructure has not been put into place. Due to large attrition rates after the completion of small projects from many donor organizations, there is a feeling of resentment and distrust towards NGOs from the recipients of these projects (Sanders and McKay 2013, 112).

MMCs are known in the NGO world as providing short-term care. Even in their title they profess to be mobile entities. An MMC's main objective is to provide temporary health care within a short period of time, then move on to other areas of need. This idea

questions both the effectiveness of health care being offered, and also, the ability to provide continuous health care for individual persons.

David Citrin, Nancy Scheper-Hughes, and Ivan Illich question the ‘medicalization’, or seemingly endless supply of drugs, which are thrown at an issue that might be more benefitted by addressing the root cause- a food crisis (Citrin 2012, 17; Illich 1975, 31; Scheper-Hughes 1992, 203). Virchow wonders whether bettering social conditions is the more important task at hand (Citrin 2012, 17)? Their argument is that MMCs provide a venue which “strip[s] the conditions that foster poverty, scarcity, sickness, and hunger of their human origins and [turn] them into problems solvable by dispensing medications or food in the short-term” (Citrin 2012, 18).

Furthermore, the term mobile medical camp might be a misnomer. Even though MMCs are mobile, many people still have to travel long distances to reach them. Results of travel times can be seen in the results section in Table 14.

Despite these critiques, NGOs and MMCs continue to exist and provide care that would otherwise not exist. “Insufficient resources, inappropriate allocation, and inadequate quality are major impediments to the delivery of effective health care that reaches the poor” (O’Donnell 2007, 2825). The problem of access to health care is two-sided: supply and demand. MMCs in Humla, Nepal, as well as the rest of the world, are striving to perfect the demand-side of health care. The reality is that many economically developing countries, including Nepal, depend on support from NGOs to provide health care for its citizens (O’Donnell 2007, 2829).

MMCs are used around the world because they have several advantages. They can provide “...autonomous functioning, rapid reacting, mobility and universality, [and]

availability...” (Glu, et al. 2012, 128). These health care facilities can provide medical care even in the most remote settings where there is a lack of health infrastructure (Glu, et al. 2012, 128). There are an estimated 1,500 MMCs in the United States alone, serving the most vulnerable population of low-income, high-risk patients. Even in the United States, these MMCs provide health care to a population that would otherwise not receive services (Hill, et al 2014, 261).

MMCs are even being used to help fight AIDS in Sub-Saharan Africa, where 3/4s of the world’s AIDS population resides. MMCs in this setting will be used to treat and track AIDS patients and rates in rural areas. They are able to provide needs assessments “...and deploy limited resources accordingly. In the future, such a network could also play a vital role in distributing antiretroviral drugs and eventually a vaccine” (Open Architecture Network 2012, 1).

The World Health Organization believes that the “lack of availability [of health care facilities and resources] is the root of the problem in many instances” (O’Donnell 2007, 2821). Due to the mountainous terrain of Humla, an infrastructure of health care buildings was either never built or has been poorly staffed, equipped, and maintained (Sanders and McKay 2013, 109). MMCs are mobile and have the ability to provide health care without a large amount of infrastructure.

MMCs have the ability to provide effective and quality health care, fulfilling requirements for the supply end of the problem of access to health care. Demand-side barriers can be addressed once supply-side barriers are controlled (O’Donnell 2007, 2820). The goal of MMCs in the long run is to provide medical treatment that is not

available through any other means, and to also increase self-sufficiency in the most vulnerable populations around the globe (Subedi 2010, 30).

Besides being able to provide health care, most NGOs provide health care for free or at major discounts. This greatly increases the amount of people who can access health care. It relieves one of the financial barriers of accessing health care for poor people. Price-elasticity of health care has a major effect on demand for the service. When health care prices are lower, the demand for health care is higher. A study by Banerjee, Duflo, Glennerster, and Kinnan in 2009 showed that most people in economically developing countries spend a large portion of their resources on obtaining health care (Dupas 2011, 2-3). In 2006, Van Doorslaer, O'Donnell, and Rannan-Eliya showed that health care expenditures further exacerbated economic conditions in already-impooverished households (Dupas 2011, 2).

MMCs may reduce morbidity and mortality and meet an unmet health need in rural communities. Specifically, MMCs provide immunizations, preventive care, and health care technologies to a population that would otherwise not receive them. Again, a major barrier to health care is the cost of the treatment. Guyatt, Ochola, and Snow have shown that most households state their main reason for not using health care treatment is because of financial costs. The use of preventative technologies by MMCs has been shown to reduce child mortality and morbidity by large numbers (Dupas 2011, 4-5).

NGOs and MMCs are filling a gap in providing health care that the government cannot or will not provide. Preventive care in economically developing countries, which is supplied through the public sector, is often sparse, not available, or not fully functioning. A study by Banerjee, Duflo, Glennerster, and Kothari, in 2010, revealed that

45% of public health care workers in Udaipur, India were absent from their stationed health posts on any given day (Dupas 2011, 5-6). Absent health care workers from health posts has been shown to reduce provider reliability and patient confidence in the quality of care. Provider reliability has been proven to increase health care accessibility. Patients will be more willing to travel if quality health care is provided. In an ideal fashion and for certain services, MMCs can provide reliable, consistent, and well-advertised health care. In Udaipur, India, child immunization rates increased from 49% in an unreliable health care setting to 78% in a reliable health care setting (Dupas 2011, 6).

MMCs provide rural areas with treatment for remedial care and information for health care prevention. "...Lack of information on illness prevention or on the effectiveness and cost-effectiveness of preventative behaviors" (Dupas 2011, 9) is a barrier to seeking health care. Numerous studies have shown that information given to households in relation to specific preventive techniques, promotion of good hygiene practices, and remedial health behavior, affects health care choices and treatment (Dupas 2011, 9-11). "Households in developing countries often lack information on the returns to investing in preventive behavior [which is unfortunate because] health behavior is quite responsive to information" if given by a reputable source (Dupas 2011, 12).

Many government-funded, public health care facilities often are not utilized because of the poor quality of health care they provide. Primary care clinics claim to provide benefits for their communities, however, a review by Filmer, Hammer, and Pritchett for the World Bank showed mixed evidence on whether primary care clinics were effective at all in their impact on population health (O'Donnell 2007, 2821). NGOs might be able to help impact population health by sending reputable and reliable MMCs

to distribute information. MMCs can remain reputable and reliable by conducting QI (quality improvement) studies. They need to be consistent and continuous in gathering data, not only pertaining to patient health, but also to patient socio-demographics and socioeconomics. It is important that patient input be gathered and taken into consideration in order to provide the best possible care.

Health related NGOs are making efforts to provide more holistic and permanent care for their recipients. The NGO that collected the data used in this thesis is working to produce better monitoring systems for patient follow-up care. Patients who need long-term care are monitored using a patient database with contact information. Follow-up testing for how well this was accomplished is currently in process (Kimber McKay, personal communication, October 2014). Compilation of this database including 1,138 participants was in response to critiques of short-term medical camps by scholars and practitioners, and to villagers' stated need for more extensive education, referral, and follow-up care.

Data Acquisition and Preparation

The data that was used for this thesis was extant data in a quality improvement (QI) study completed by Nepali citizens working for an NGO. I was commissioned to help analyze the existing data from this respective NGO. I was handed this de-identified data and did not collect this data. Therefore, the data gathered did not undergo IRB review or approval.

Two Nepali doctors collected and entered data from 1,138 participants who attended one of six MMCs in Humla district of Northwestern Nepal in December of 2013. All patients attending the MMC were seeking remedial health care. Remedial

health care restores the health of the patient when there is an illness, as opposed to preventive health care, which reduces the risk of bad health and illness (Dupas 2011. 7).

Surveys were given to all patients. If the patient was a child the survey was given to their parent. The information collected in this case included the child's reason for seeking treatment but the socioeconomic measures, socio-demographic measures, health care measures, and measures of health care accessibility of the parent.

The data were collected in the form of hard copy for every patient who attended any of six MMCs and entered into the Nepali language. There was some translation from Tibetan into Nepali in the Buddhist villages by members of NGO camp staff, but no English translation occurred in the field. Locations of data collection were six sites within the Humla district of Nepal: Chauganphaya, Dandaphaya, Kermi, Muchu, Syada, and Yalbang.

After the camp, the physical data were translated into English and Excel, at the same time. The data collectors were literate villagers with high school educations and the data were digitized by the chair of the Department of Social Work at Classic College in Kathmandu, Nepal. I received the database in Excel format and transferred the data into SPSS format, concurrent with the organization needed for the respective analyses. Myself, a second research assistant, and the NGO's director of research performed quality checks in Excel and SPSS.

Limitations in Data Acquisition

Data limitations were present because data were based upon survey data collected by other workers from an NGO. In order to produce a fuller examination of socioeconomic influences on health care seeking behavior, additional data would have

been required. Specific details about the severity of patient medical conditions could provide data for fuller SES analyses. Furthermore, qualitative data for variables analyzed with SES, such as healthcare indicators (amount of time since previous treatment was received, type of health care facility visited for previous treatment, location of previously received treatment, person who provided work replacement while patient was seeking health care, whether children were brought with patient or left at home, person who provided childcare while patient was seeking health care) would help provide a fuller and more robust analysis. However, because the dataset used for this thesis was given to me with set variables, I conducted an analysis of socioeconomic influences within the given dataset.

I would also like to acknowledge the people who were too sick to walk to an MMC to receive health care. This dataset did not include the sickest people who were unable to travel by foot to an MMC or the poorest people, who could not afford to lose work by taking time to walk to the MMC and receive health care.

Analysis Methodology

This thesis included two sub-projects. The first project used SPSS (Statistical Package for the Social Sciences) to produce both descriptive and inferential analyses. This project's goal was to assess the contribution of SES to health care seeking behavior. This project tested the hypothesis that SES has an effect on one's health care seeking behavior. Variables tested by analyses in SPSS included SES indicators, socio-demographic indicators, health care access indicators, and sickness indicators.

Socioeconomic status (SES) indicators: Socioeconomic measures were gotten from each patient based upon a threefold (high/medium/low) self-identifying method. Levels

of SES are identified by the amount of land one has, the amount of animals one has, and food security (Humla District Development Committee 2000, 18-19). Humla district is a primarily egalitarian. Eighty-one percent of households are income dependent on agriculture and 12% are income dependent on wage labor (Humla District Development Committee 2000, 18). This rapid appraisal method of identifying SES is an anthropological method “which uses qualitative measures of SES and is designed to rank or stratify participants in wealth categories” (van Bodegom et al. 2009, 69). This method has shown to be accurate in representing one’s actual SES through comparison with other forms of SES-gathering methodology, including the Demographic Health Survey (DHS) wealth index and interview-reported wealth (van Bodegom et al. 2009, 67).

Socio-demographic indicators: Socio-demographic indicators were compared against SES to ascertain any relationships. Socio-demographic indicators included age, gender, marital status (single, monogamous, polygynous, polyandrous (a marriage system in which a woman marries a set of brothers), monogamous but used to be polyandrous), number of adults living in household, village of residence, and village where respondent was born.

Health care access indicators: Variables were measured which indicated one’s level of accessibility to health care. The amount of time traveled to receive health care (in minutes) was measured. All patients seen at one of six MMCs traveled by foot to receive health care. Trails in Humla district are not built for transportation other than walking, and villagers are too poor for other forms of transportation. The amount of time since previous treatment was received (in months), type of health care facility visited for previous treatment (e.g., none, health post, health camp, hospital, traditional,

unspecified), and location of previously received treatment were measured. The person who provided work replacement while patient was seeking health care (e.g., no one, self, spouse, affine (non-blood relation), consanguine (blood relation), non-relatives, other family members, child went to school) was also provided. Work replacement was provided for everyone, regardless of whether they had to travel from another village or received health care at an MMC in their same village of residence. Whether children were brought with patient or left at home was measured, the person who provided childcare while patient was seeking health care (e.g., no one, self, spouse, affine, consanguine, non-relatives, other family members, child went to school) was measured, and village where health care was sought was measured.

Sickness indicators: One sickness indicator was measured and compared with SES in search of a relationship. This variable included chief medical complaint/ reason for attending MMC (e.g., fever, head pain, body pain, respiratory, vision impairment, GI tract, genital, reproductive or STI related, tooth or mouth pain, ear ache or hearing impairment, allergy or itching, wound, other, which included dizziness, shivering, heart pain, worms, nose pain, throat pain).

Descriptive Statistical Analysis

The analyses conducted were based upon nominal and ratio data. I performed frequency analyses for each nominal variable being tested: SES, gender, marital status, village of residence, village where respondent was born, type of health care facility visited for previous treatment, location of previously received treatment, person who provided work replacement while patient was seeking health care, whether children were brought with patient or left at home, person who provided childcare while patient was

seeking health care, village where health care was sought, and chief medical complaint/ reason for attending MMC. I performed descriptive statistical analyses for all ratio variables being tested: age, number of adults living in household, amount of time traveled to receive health care, and amount of time since previous treatment was received.

Prior to the inferential analysis performed, I conducted tests upon the data to investigate their quality. I produced a test for significant outliers. I performed a distribution analysis on all relevant variables by finding the kurtosis and skewness. Results from this test influenced the use of either parametric "...statistical procedures that rely on assumptions about the shape or form of the probability distribution from which the data were drawn" or non-parametric parametric analyses "...statistical procedures that do not rely on assumptions about the shape or form of the probability distribution from which the data were drawn" (Hoskin N.D., 1). To produce more robust results with the highest statistical power, I hoped that results would be normal and parametric tests could be performed (Garth 2008, 4-5). Prior to an analysis of comparison of means, I performed Levene's test for homogeneity of variances to assess the assumption that the populations of samples were equal.

Inferential Statistical Analysis

I conducted two types of inferential analyses: one-way ANOVA (analysis of variance) and chi-square test. I conducted a one-way ANOVA analysis for both socio-demographic variables, SES, sickness indicators, and health care access indicators (Aerd 2013, 1). I performed analyses to test whether SES determined the number of adults living in the patient's household, whether SES determined if and when previous treatment was sought for remedial illnesses, and if SES determined the amount of time

patients traveled to receive health care. I conducted a one-way ANOVA analysis after assumptions in my data were validated (Aerd 2013, 1). The dependent variables (number of adults living in household, amount of time since previous treatment was received, and travel time needed to reach MMC) were continuous and measured at the ratio level. The independent variable (SES) consisted of three categorical and independent groups: low, medium, and high. There was no relationship between the independent variables, either within or between groups. These results allowed for independence of observations (Aerd 2013, 1).

I conducted a chi-square test to determine whether there was a statistically significant association between SES and nominal variables, such as socio-demographic indicators, health indicators, and health care access indicators. Socio-demographic indicators included village where respondent was born and village of residence, patient gender, and marital status. Health indicators included chief medical complaint/ reasons for attending MMC. Health care access indicators included type and place of treatment already received, work replacement and childcare, and respective MMC where patient received health care.

Spatial Examination of Health Care Seeking Routes

In the second project, I used ArcGIS (geographic information system) and Google Earth to produce a map that showed walking routes between GPS-located patient villages and MMCs. This map calculated the distance that was needed to travel between villages and MMCs. For each route I created a distance and elevation profile.

From the dataset, I matched participants from their village of residence to the respective MMC they attended. Using aerial photographs from Google Earth, I located

the six MMCs and all villages of residence. I mapped the GPS locations using ArcGIS. Then I mapped the most commonly used routes from all villages to the six MMCs using Google Earth route. The ending result included routes from GPS locations of six MMCs and 14 villages.

The goal of this project was to show the distance, as well as gain and loss in elevation, required to access health care for villagers within Humla, Nepal. This route map could be used against the MMC database in measuring costs of travel to determine distance decay (the decrease in utilization with distance) (Mueller et al. 1998, 878).

Results of Statistical Analyses and Spatial Construction of Routes

Table 1 shows the results of a frequency analysis of chief medical complaint at all MMCs for both females and males. Although there were only 1138 patients, many patients reported multiple injuries and medical complaints. Body pain in both males and females was the highest medical complaint.

Table 1: Patient Chief Medical Complaint

Medical Complaint	Patient Count	Percentage	Female	Percentage	Male	Percentage
Body Pain	511	28%	339	28%	172	29%
Respiratory	294	16%	182	15%	112	19%
GI Tract	286	16%	212	17%	74	12%
Head Pain	223	12%	170	14%	53	9%
Vision Impairment	191	11%	123	10%	68	11%
Other	172	9%	102	8%	70	12%
Mouth Pain	66	4%	48	4%	18	3%
Ear Pain/ Hearing Impairment	46	3%	19	2%	27	5%
Genital, STI, Reproductive	24	1%	24	2%	0	0%
Total	1813	100%	1219	100%	594	100%

Table 2 shows the results of a frequency analysis of patient SES for both females and males. Although there were 1138 patients, some did not provide a self-report of SES. The highest category of SES was the middle level, which indicated that most patients considered themselves not to be rich or poor, but somewhere in between.

Table 2: Patient Socioeconomic Status

Level of SES	Patient	Percentage	Female	Percentage	Male	Percentage
Low	174	15.7%	116	10.5%	58	5.2%
Middle	900	81.3%	602	54.4%	298	26.9%
High	33	3%	26	2.3%	7	0.6%
Total	1109	100%	744	67.2%	363	32.8%

Tables 3a and 3b show the results of a frequency analysis of patient gender and age. Results conclude that more female than male patients attended MMCs. The highest age group range to seek health care at an MMC was from ages 21 to 30.

Table 3a: Patient Gender and Age in Years (Age 1-50)

Gender	Age									
	1-10	Percent	11-20	Percent	21-30	Percent	31-40	Percent	41-50	Percent
Female	3	43%	99	70%	202	79%	175	76%	109	62%
Male	4	57%	42	30%	53	21%	54	24%	68	38%
Total	7	100%	141	100%	255	100%	229	100%	177	100%

Table 3b: Patient Gender and Age in Years (Age 51-90)

Gender	Age									
	51-60	Percent	61-70	Percent	71-80	Percent	81-90	Percent	Total	Percent
Female	96	56%	58	54%	13	41%	3	38%	758	67%
Male	75	44%	49	46%	19	59%	5	62%	369	33%
Total	171	100%	107	100%	32	100%	8	100%	1127	100%

Table 4 shows the results of a frequency analysis of patient SES and respective MMC attended. The MMC at Chauganphaya treated more patients than any other MMC.

Table 4: Patient Socioeconomic Status and MMC Attended

MMC Attended	SES									
	Low	Percentage	Medium	Percentage	High	Percentage	Total	Percentage		
Chauganphaya	91	52%	366	41%	6	18%	463	42%		
Dandaphaya	58	33%	188	21%	7	19%	253	23%		
Kermi	0	0%	54	6%	0	0%	54	5%		
Muchu	1	0%	47	5%	1	3%	49	4%		
Syada	15	9%	186	21%	18	55%	219	20%		
Yalbang	8	6%	57	6%	1	3%	66	6%		
Total	174	100%	902	100%	33	100%	1109	100%		

Tables 5-10 show the results of a frequency analysis for each MMC and the villages of residence that most often frequented the MMC. These results allowed us to

better understand the routes that patients traveled to seek health care. Results from this frequency analysis were indicative of proper placement of MMC locations for residents of the Humla district. Results showed that the MMC at Chauganphaya was most frequented by villagers living in Chauganphaya, the MMC at Dandaphaya was most frequented by villagers living in Dandaphaya, the MMC at Kermi was most frequented by villagers living in Kermi, the MMC at Muchu was most frequented by villagers living in Muchu, the MMC at Syada was most frequented by villagers living in Syada, and the MMC at Yalbang was most frequented by villagers living in Yalbang. These results were expected. If results showed that MMCs were not providing the largest percentage of their health care to residents within their own village, this would be indicative of providing a misplaced MMC. These results acknowledge proper placement of MMCs provided by the NGO in the Humla district.

Table 5: MMC (Chauganphaya) Attendance Based Upon Village of Patients' Residence

Village of Residence	Number of Patients	Percentage
Chauganphaya	127	27%
Kholsi	82	17%
Dharapori	62	13%
Hepka	61	13%
Dinga	48	10%
Pamlatum	32	7%
Gadapori	20	4%
Syada	18	3%
Other	27	6%
Total	477	100%

Table 6: MMC (Dandaphaya) Attendance Based Upon Village of Patients' Residence

Village of Residence	Number of Patients	Percentage
Dandaphaya	226	88%
Tulin	15	6%
Dharapori	6	2%
Chauganphaya	4	2%
Other	5	2%
Total	256	100%

Table 7: MMC (Kermi) Attendance Based Upon Village of Patients' Residence

Village of Residence	Number of Patients	Percentage
----------------------	--------------------	------------

Kermi	52	96%
Chyaduk	2	4%
Total	54	100%

Table 8: MMC (Muchu) Attendance Based Upon Village of Patients' Residence

Village of Residence	Number of Patients	Percentage
Muchu	30	60%
Dumkot	11	22%
Yari	8	16%
Thawang	1	2%
Total	50	100%

Table 9: MMC (Syada) Attendance Based Upon Village of Patients' Residence

Village of Residence	Number of Patients	Percentage
Syada	194	85%
Sata	13	6%
Kholsi	12	6%
Yangu	5	2%
Other	4	1%
Total	228	100%

Table 10: MMC (Yalbang) Attendance Based Upon Village of Patients' Residence

Village of Residence	Number of Patients	Percentage
Yalbang	34	49%
Yangu	20	29%
Chala	16	22%
Total	70	100%

Table 11 shows a frequency analysis of childcare decision making for patients who traveled from their village of residence to one of six MMCs to receive health care. Patients were asked whether they brought their children or left them at home. Results showed that most parents left their children at home.

Table 11: Patient Childcare Decision Making

	Frequency	Percentage
Left Children at Home	757	70.7%
Brought Children	312	29.2%

Table 12 shows a frequency analysis of childcare decision making for patients who traveled from their home village to one of six MMCs to receive health care. Out of the patients who left their children at home (see Table 11) patients were asked to give their childcare provider (given in the form of the relationship to health care seeking

parent). Options included no one, spouse, affine, consanguine, and other (unspecified).

Results showed that most parents left their children at home with no one watching them.

Table 12: Patient Childcare Provider

	Frequency	Percentage
No one	416	52.2%
Spouse	235	29.5%
Affine	31	3.9%
Consanguine	99	12.4%
Other	28	3.5%

* Multiple answers could have been given, so numbers don't add up to equal 757 (Table 11, Left Children at Home)

Table 13 shows a frequency analysis of work replacement decision making for patients who traveled from their home village to one of six MMCs to receive health care. Patients were asked to give their work replacement provider (given in the form of the relationship to patient). Options included no one, spouse, affine, consanguine, and other (unspecified). Results showed that the patient's spouse usually provided work replacement for the patient.

Table 13: Patient Work Replacement

	Frequency	Percentage
No one	165	14.8%
Spouse	517	46.2%
Affine	182	16.3%
Consanguine	383	34.3%
Other	8	.7%

Table 14 shows the descriptive statistics for amount of time traveled to MMC (in minutes) for 1126 patients who traveled from their home village to one of six MMCs to receive health care. The results excluded 1 major outlier. Results indicated that the mean amount of time traveled to MMC was 40.9 minutes, the maximum was 720, and the minimum was one minute.

Table 14: Amount of Time Traveled to MMC (in minutes), excluding outliers

Valid	1126
Missing	16
Mean	40.9423

Median	15
Mode	5
Std. Deviation	66.3517
Minimum	1
Maximum	720

In order to conduct an ANOVA analysis, I produced a test for significant outliers on all dependent variables so I could extract any potential outliers that could have a negative effect on the ANOVA analysis. Without outliers the validity of the results would be maintained. Dependent variables tested for significant outliers included the number of adults living within the respondent's household, when previous treatment was received (in months), and amount of time required to travel to the MMC (in minutes). Figures 2- 4 show results. There were no outliers within the three dependent variables.




Figure 2: Test for Outliers within Number of Adults in Household




Figure 3: Test for Outliers within Amount of Time Traveled to MMC (in minutes)




Figure 4: Test for Outliers within When Previous Treatment was Received (in months)

I conducted a test for normal distribution on the dependent variables, based upon the assumption that an ANOVA analysis needs to be performed on a normally distributed dataset. I conducted a test to determine the skewness and kurtosis of distribution on all three dependent variables used in the one-way ANOVA analysis. Dependent variables included the number of adults living within the respondent’s household, when previous treatment was received, and amount of time traveled to MMC.

Table 15 shows results. The skewness (which signifies how symmetrical the distribution is) must equal 0 to be symmetrical and have a normal distribution. These values are shown below to be substantially asymmetrical. The values have a positive skew, which means the distribution is asymmetrical with a long tail to the right. The kurtosis values are well above the expected kurtosis of 0, which means they have a positive kurtosis. Although the dependent variables were not shown to be normally distributed, my decision to use a parametric test was influenced by the sort of data being analyzed. Even though these distributions didn’t seem to be normal and an assumption was violated, the ANOVA test still seemed the most appropriate test used to analyze a ratio variable against a nominal variable.

Table 15: Test for Normal Distribution of Dependent Variables

	Skewness	Kurtosis
Number of Adults in Household (above age 18)	1.541	6.083
When Previous Treatment was Received (in months)	33.044	1103.279
Amount of Time Traveled to MMC (in minutes)	2.074	4.019

Table 16 shows the results from Levene’s test for homogeneity of variances for SES and all dependent variables tested within the ANOVA analysis. For all variables except the number of adults in household sig > .05, which meant that the obtained differences in these sample variances were not significantly different. For variables with

a significance level greater than .05, the null hypothesis of equal variances could not be rejected. I therefore concluded that there was no difference between the variances in the population. Because of this parametric tests could be performed on the samples.

However, because the significance level was lower than .05 within the number of adults in household, the null hypothesis had to be rejected. This meant that a one-way ANOVA analysis could not be performed on the number of adults in each household because the obtained differences in the sample of variances was significantly different.

Table 16: Homogeneity of Variances for SES

	Levene Stat	Sig
Number of Adults in Household (above age 18)	3.569	.029
When Previous Treatment was Received (in months)	1.794	.168
Amount of Time Traveled to MMC (in minutes)	.182	.834

Table 17 shows the results of the descriptive statistics for a one-way ANOVA.

This analysis tested whether number of adults living in household, when previous treatment was received, and amount of time traveled to MMC differed based upon level of SES. These three tests were included because I was expecting people with a higher SES to have a lower number of adults living in each household, have a greater frequency of treatment, lower amount of time since last treatment was received, and have a lower amount of time traveling to MMCs due to greater options for transportation.

Table 17: Descriptive Statistics of one-way ANOVA for SES

Independent Variables	SES		
	Low	Medium	High
Number of Adults in Household	Mean: 3.2	Mean: 3.8	Mean: 3.3
	Minimum: 0	Minimum: 0	Minimum: 0
	Maximum: 8	Maximum: 20	Maximum: 9
When Treatment was Received (in months)	Mean: 13	Mean: 8.4	Mean: 3.9
	Minimum: .1	Minimum: 1	Minimum: .1
	Maximum: 120	Maximum: 180	Maximum: 12
Amount of Time Traveled to MMC (minutes)	Mean: 45	Mean: 64	Mean: 45.2
	Minimum: 1	Minimum: 1	Minimum: 2
	Maximum: 720	Maximum: 1500	Maximum: 300

Table 18 shows the results of a one-way ANOVA analysis for SES and all dependent variables (except for number of adults in household, which could not be analyzed using an ANOVA analysis because it did not pass Levene’s test for homogeneity of variances). The test indicated that there were no statistically significant differences in the mean of when treatment was received and amount of time traveled to MMC. Both significance levels were greater than 0.05. I expected that people with higher SES would be more likely to seek treatment more frequently. The finding suggested that, irrespective of wealth status, there was no difference in the frequency of treatment sought. Similarly, I expected that people with higher SES would be more likely to spend less time traveling to an MMC, as they might be able to afford transportation services other than walking. However, again, irrespective of wealth status, there was no difference in the amount of time required to travel to an MMC.

Table 18: Results for One-way ANOVA Analysis for SES and Dependent Variables Tested

Dependent Variable	Sig
When previous treatment was last obtained	.214
Amount of time required to travel to MMC	.932

I performed multiple Pearson chi-squared analyses on the SES of patients to test for an association between SES and other variables. Analyses tested the null hypotheses that SES would not be related to or have a relationship with any of the following variables: marital status, patient gender, chief medical complaint, childcare decision making, work replacement, type and place of previous treatment received, respondent’s village of birth, residence, and which MMC the patient attended. Table 19 shows the results of statistically significant variables that were shown to have a relationship with SES.

I was curious to see whether there would be a relationship between SES and people of differing marital statuses or chief medical complaint. I would have expected to see a relationship between SES and patient gender, as females in most of Nepal generally have a lower SES than males. Results showed no statistically significant relationship between SES and marital status ($p = .796$), patient gender ($p = .355$), or chief medical complaint ($p = .088$).

I would have expected to see an relationship between SES and childcare decision-making. I predicted that people with a higher SES would be more able to attend an MMC because it would be easier to find or hire childcare with a higher SES. I tested the null hypothesis that SES and the decision to leave children at home or bring them to the MMC were not related using a chi-squared analysis. The hypothesis was rejected because $p < .05$. However, I also tested the relationship between SES and the provider of childcare using a chi-squared analysis. This relationship was not significant ($p > .05$). A relationship was proven between SES and whether the patient brought their children to the MMC or left them at home. However, there was not a relationship between the provider of childcare (spouse, affine, consanguine, non-relatives, and other unspecified family members) and SES.

I expected to find relationship between SES and work replacement. I assumed that people with a higher SES could afford to find or hire work replacement easier than people with a lower SES. The results, shown in Table 19, provide evidence for a relationship between SES and work replacement.

I expected to find a relationship between SES and type and place of previous treatment received. I assumed that people with a higher SES would have traveled further

to seek medical treatment, and would also have sought westernized treatment, due to higher levels of education. I performed a chi-squared analysis between SES and type of treatment already received. The results ($p=.266$) lead to the conclusion that no evidence could be found for an association between SES and type of treatment already received. However, if the patient had already received treatment, the location treatment was received at was shown to have a relationship with SES.

I was curious to see whether there was a relationship between SES and respondent's village of birth, residence, and MMC the patient attended. I did not know whether villages in the Humla district all had similar or different SES. Varied SES between villages could have an affect on health care seeking behavior throughout the district. I performed a chi-squared analysis between SES and village where respondent was born, village of residence, and village in which the patient attended the MMC. A relationship was shown between SES and village where respondent was born, village of residence, and village where information was collected.

Table 19: Chi-Square Test between SES and Variables

Variable	Sig.	df	Number of cases
Decision to Bring Children or Leave them at Home	.021	2	1048
Location of Treatment Previously Received	.023	62	404
Village where Respondent was Born	.000	122	1102
Village of Residence	.000	60	1109
Village where Information was Collected	.000	16	1109
Work Replacement	.034	18	1097

Project II involved locating the villages of patients and MMCs which patients attended in Humla, Nepal. Figure 5 shows a map of the most-used routes that were traveled from patient villages to MMCs.




Figure 5: Walking routes from villages to MMCs (Google earth 2014)

Discussion

In this project, I attempted to identify barriers that prevented patients from seeking health care within Humla, Nepal. The analyses I conducted helped to bring a better understanding of socioeconomic determinants on health care seeking behavior from six MMCs. I used the data collected from an NGO to help identify specific barriers, and to shed light on their relative importance to health care seeking behavior among local people. I conducted a one-way ANOVA analysis to evaluate the influence of socioeconomic barriers on health care seeking behavior. I conducted multiple chi-squared analyses to identify relationships between SES and health care seeking behavior. I produced a map to help locate villages and MMCs. From GPS marked locations, I created

routes to help NGOs better understand the cost of travel associated with distance needed to walk and elevation gains and losses.

This thesis tested my hypothesis that SES effects health care seeking behavior. To test this hypothesis, I conducted a one-way ANOVA analysis, which found that SES affected none of the variables tested. The one-way ANOVA analysis tested the effect of SES on amount of time since treatment was last received and amount of time that was required to travel to MMC. The one-way ANOVA analysis that tested SES and amount of time since treatment was last received was relevant in its findings, in that, irrespective of wealth, people were not more likely or less likely to seek health care treatment more frequently. Since this finding seems surprising, future tests could more extensively explore the relationship between SES and frequency of health care seeking behavior. Reasoning for this could be explained utilizing a qualitative search for meaning and reasoning.

The results of the one-way ANOVA analysis that tested the effect of SES on amount of time required to travel to MMC are encouraging. These results have profound effects on the equality of health care treatment being provided to the villagers in the Humla district. These results mean that both rich and poor people are as likely to spend the same amount of time traveling to an MMC. Generally, wealthier people have better access to health care. These results indicate that, at least geographically, there is no advantage in being wealthy and accessing health care than being poor in Humla.

Multiple chi-squared analyses determined there was a relationship between SES and the decision to bring children or leave them at home, provision of work replacement, location of treatment previously received, village where respondent was born, village of

residence, and village where information was collected. SES had an effect on the decision to bring children or leave them at home and on work replacement. Perhaps people with a higher SES are better able to find or hire childcare and work replacement. Perhaps one's SES has an effect on the strength of one's social network and support system. Although it is interesting to find out that there is a positive relationship between SES and the decision to bring children or leave them at home and work replacement, further analyses need to be conducted in order to find out why this is happening. What is the relationship between SES and childcare and SES and work replacement? Once the relationship is better defined, qualitative analysis can help to provide in-depth reasoning.

A positive relationship between SES and location of previous treatment received was proven significant. This is interesting, considering a one-way ANOVA analysis showed that amount of time since previous treatment was received and SES did not have a relationship. This means SES does not have an effect on the frequency of treatment received, but it does have an effect on where treatment is received.

Village where respondent was born, village of residence, and village where information was collected were all shown to have a positive relationship with SES. This implies that the village that people reside in has an impact on their SES. Further exploration needs to find out explanatory reasons for this relationship. Perhaps some villages are wealthier than others. This finding could have an impact on health care seeking behavior. This research has helped to better understand which variables SES affects, however, continuing research needs to be conducted in order to understand how and why these variables are associated with SES.

Suggestions for Further Research

Suggestions for further research include using the route map from villages to MMCs within Humla, Nepal to conduct spatial analyses to determine distance-related health seeking behavior barriers, known as distance decay. Calculating the distance and elevation gain and loss of each route to determine whether or not the combined factors have any effect on the willingness for patients to travel to seek health care could do this. This route map could be used against the MMC database in measuring costs of travel to determine distance decay (the decrease in utilization with distance) (Mueller et al. 1998, 878).

Research is needed in order to address the duplication of efforts NGOs are providing. Further research includes identifying areas in Humla, Nepal where services are being provided, what kinds of services are being provided, and what NGO or organization is providing those services. The goal of this research would be to combine efforts of various NGOs to eliminate duplication of efforts and increase the efficiency of services being offered.

Research should also be conducted in relation gender disparities in seeking health care at MMCs in Humla district. In the results section, 758 females were identified as seeking health care from an MMC, and only 369 males were identified as seeking health care from an MMC. This means that 67% of patients seen at an MMC were females and only 33% were males. Further research needs to identify the reasoning behind this disparity. Perhaps this number is equivalent to the amount of females and males living in the villages due to males traveling out of the region for work. Perhaps this number is indicative of females receiving health care when they take their children to receive health

care. Further research should identify the reasoning for this discrepancy in order to better provide health care to males within Humla, Nepal.

Implications for MMCs in Health Care Behavior

Access to health care is a problem for many people in rural areas and in economically developing countries (O'Donnell 2007, 2830). Effective health care is oftentimes not available through the state, so NGOs have taken up a large part in trying to provide health care for this ostracized group (Sanders and McKay 2013, 109). In Humla, Nepal, one such intervention has been applied through the use of MMCs. Although there is much criticism surrounding MMCs and the work of NGOs in economically developing countries and regions such as Humla, Nepal, NGOs do provide a service to people that could not and would not be provided without these organizations.

Strategies are needed in order to raise utilization of health care in the Humla district of Nepal. MMCs can help to reduce barriers from both the demand and supply-side. From a demand-side, MMCs can provide reliable, reputable, and quality health care. From a supply-side NGOs can help to inform community households and individuals about the positive effects of both preventative and remedial health care treatments. They can reduce the distance that patients need to travel in order to receive health care by providing MMCs closer to rural villages. They can provide demand-side economic incentives (O'Donnell 2007, 2831) to offset the socioeconomic cost of health care by providing food or money incentives worth the amount of work missed. NGOs can use MMCs in continuing to provide social impact analyses on rural patients with the goal of furthering the utilization of health care to this group of people. "The impact on the

average utilization of health services should be identified and differential effects with socioeconomic characteristics tested” (O’Donnell 2007, 2831).

This research drew upon the theoretical framework of critical medical anthropology to study the social production of illness in order to help better understand socioeconomic factors imbedded in health care seeking behavior. This study was important in the continuation and dedication of social scientists to study societies and the greater needs of those societies before introducing social changes. It drew upon the health behavioral model in a focus specifically on the socioeconomic determinants of health care seeking behavior.

Access to health care is influenced by cultural and historical conditions, as well as hegemonic systems of power (Alexandrakis 2001, 75). This project helped bring knowledge and better understanding of barriers to health care. I conducted these analyses in hopes of influencing policy and practice relating to making health care accessible even to the most remote-dwelling people, using the most flexible form of health care delivery – the mobile medical camp. The framework provided by social psychology’s health behavior model regarding perceived barriers influenced my thinking and analysis of my findings. As perceived barriers are noted by Hyder and Morrow to be the most predictive factor in influencing health care seeking behavior (2005, 51), this model was used in measuring social costs of accessing health care.

References

- Action Contre La Faim (ACF)
2006 Nutritional Exploratory Mission: Humla and Mugu Districts: Nepal.
- Adara Development- Bridging Worlds.
<http://www.adaragroup.org/foundation-projects/nepal/>, accessed November 11, 2014.
- Aerd Statistics
2013 One-way ANOVA in SPSS. Lund Research Ltd. <https://statistics.laerd.com/spss-tutorials/one-way-anova-using-spss-statistics.php>, accessed January 21, 2015.
- Alexandrakis Othon
2011 Medical Anthropology: The Development of the Field. Totem: The University of Western Ontario Journal of Anthropology. 9(1):73-77.
- Bakeera, Wamala, Galea, State, Peterson, Pariyo
2009 Community perceptions and factors influencing utilization of health services in Uganda. International Journal for Equity in Health. 8(25):1-12.
- Balabanova, Dina, Parkhurst, Justin, McKee, Martin, McPake, Barbara
2006 Access to health care: taking into account health systems complexity. UK: Health Systems Development.
- Basnett, Yurendra
2009 From Politicization of Grievances to Political Violence: An Analysis of the Maoist Movement in Nepal. Dissertation. Development Studies Institute. London.
- Bedford, Juliet
2012 Qualitative study to identify solutions to local barriers to care-seeking and treatment for diarrhoea, malaria and pneumonia in select high burden countries: Report on findings from Kenya. New York: UNICEF.
- Bork-Huffer, Tabea
2012 Migrants' Health Seeking Actions in Guangzhou, China: Individual Action, Structure and Agency: Linkages and Change. Stuttgart. Franz Steiner Verlag.
- Central Bureau of Statistics
2012 National Population and Housing Census 2011. Government of Nepal. Kathmandu, Nepal
- Citrin, David M.
2010 The Anatomy of Ephemeral Health Care: "Health Camps" and Short-Term Medical Volontourism in Remote Nepal. Studies in Nepali History and Society 15(1):27-72.

- Citrin, David M.
2012 The Anatomy of Ephemeral Care: Health, Hunger, and Short-Term Humanitarian Intervention in Northwest Nepal. PhD Dissertation, Department of Philosophy, University of Washington.
- Citrin, David M.
2011 “Paul Farmer Made Me Do It”: A Qualitative Study of Short-Term Medical Volunteer Work in Remote Nepal. MA Thesis, Department of Public Health, University of Washington.
- Coreil, Jeannine
2008 Behavior and Social Science Theory. In Theoretical Foundations.
- Creel, Sass, Yinger
1996 Client-Centered Quality: Clients' Perspectives and Barriers to Receiving Care. New Perspectives on Quality of Care: No 2. Population Council and Population Reference Bureau.
- Daniel, Tsufit, Hashman, Nadia, Lanning, Kiryn, Shintani, Mai, Yadav, Vivek
2012 Health is Wealth : Health Care Access for Dalit communities in Saptari, Nepal. Samata Foundation.
- De los Angeles Nunez Carrasco, Lorena
2008 Chapter II: Theoretical Approaches and Key Concepts in Medical Anthropology (p40-61). In Living on the margins: illness and health care among Peruvian migrants in Chile. PhD thesis. Leiden University.
- Deogaonkar, Milind
2004 Socio-economic inequality and its effect on health care delivery in India: Inequality and health care. Electronic Journal of Sociology. ISSN: 1198 3655.
- Digital Himalaya's Nepal Maps Collection
2014 Humla District. Digital Himalaya.
<http://www.digitalhimalaya.com/collections/maps/nepalmaps/>, accessed March 10, 2015.
- District Development Committee (DDC)
2004 District Profile of Humla. Humla, Nepal DDC.
- DFID Nepal
2008 <https://www.gov.uk/government/world/organisations/dfid-nepal>, accessed November 11, 2014.
- Dupas, Pascaline

- 2011 Health Behavior in Developing Countries. UCLA & NBER. Annual Review of Economics Vol:3.
- Ensor, Tim, Cooper, Stephanie
2004 Overcoming Barriers to Health Service Access and Influencing the Demand-side Through Purchasing. The World Bank.
- Farmer, Paul
1999 Infections and Inequalities: The Modern Plagues, updated edition. Berkeley: University of California Press.
- Ferguson and Gupta
2002 Spatializing States: Toward an Ethnography of Neoliberal Governmentality. *American Ethnologist* 29(4): 981-1002.
- Fricke, T.E.; Axinn, W.G.; Thornton, A.
1993 Marriage, social inequality, and women's contact with their natal family in alliance societies—Two Tamang examples, *American Anthropologist*, 95(2):395-419.
- Garth, Andrew
2008 Analysing data using SPSS (A practical guide for those unfortunate enough to have to actually do it.) Sheffield Hallam University.
- Glu, Kucher, Ziniukov, Terliuk, Maksimets, Zalesskii
2012 The mobile medical clinic: a role of mobile physician crews in the provision of health care for rural populations. *Lik Sprava* 1-2:128-32.
- Grocke, Michelle
2013 Doctoral Dissertation Research: How New Chinese Commercial Food Channels Alter the Nutrition, Food Security, and Psychological Well-Being of Himalayan Villagers. In progress.
- Gupta, Anirudha
1964 Politics in Nepal: A Study of Post-Rana Political Developments and Party Politics. Bombay: Allied Publishers Private Ltd.
- Hausmann-Muela, Susanna, Ribera, Joan Muela, Nyamongo, Issac
2003 Health-seeking behavior and the health system response. DCPP Working Paper No. 14.
- HEAD Nepal
2013 <http://headnepal.org/abouthumla.html>. Accessed on November 11, 2014.
- Hill, Caterina F., Powers, Brian W., Jain, Sachin H., Bennet, Jennifer, Vavasis, Anthony, Oriol, Nancy E.

- 2014 Mobile Health Clinics in the Era of Reform. *American Journal Managed Care*. 20(3): 261-264.
- Himalayan Children Society
2009 www.himalayanchildrensociety.org. Accessed on November 11, 2014.
- Humla District Development Committee
2000 Baseline Survey Report: Humla District. Decentralised Planning for the Child Programme. http://un.org.np/sites/default/files/report/tid_188/Humla-Report.pdf. Accessed on May 01, 2015.
- Hyder, Adnan Ali, Morrow, Richard H.
2005 Culture, behavior, and Health (pg 41-66). In
- Illich, Ivan
1975 *Medical Nemesis: The Expropriation of Health*. Toronto: Bantam Books.
- Jacobs, Bart, Ir, Por, Bigdeli, Maryam, Annear, Peter Leslie, Damme, Wim Van
2012 Addressing access barriers to health services: an analytical framework for selecting appropriate interventions in low-income Asian countries. *Health Policy and Planning* 27:288-300.
- Justice, Judith
1999 Neglect of Cultural Knowledge in Health Planning: Nepal's Assistant Nurse-Midwife Program. R.A. Hahn (ed.) *Anthropology in Public Health: Bridging Differences in Culture and Society*. Oxford: Oxford University Press, Inc. 327-344.
- Justice, Judith
1986 *Policies, Plans, & People: Foreign Aid and Health Development*. Berkeley and Los Angeles, California: University of California Press.
- Kroeger, Axel
1983 Anthropological and Socio-Medical Health Care Research in Developing Countries. *Soc. Sci. Med.*17(3):147-161.
- LaFond, Anne
1995 *Sustaining Primary Health Care*. New York: The Save the Children Fund.
- MacKian, Sara
2003 A Review of Health Seeking Behavior: Problems and Prospects. *Health Systems Development Programme*.
- McElroy, A.
2002 Medical Anthropology. In D. Levinson, M. Ember's *Encyclopedia of Cultural Anthropology*. Henry Holt. New York 1996.

- McKay, Kimber
2002 Health Needs in Two Ethnic Communities of Northwestern Nepal. *Contributions to Nepalese Studies* 29(2): 241-273.
- McKay, Kimber H.
2003 Challenges to health care access in Maoist Nepal. *Himalaya*, XXIII(2): 43-46.
- McKay K. H., A. Zhand, C. Sanders, and G. Nepali
2007 Responses to Innovation in an Insecure Environment in Rural Nepal. *Mountain Research and Development* 27(4): 302–307.
- Muller, Ivo, Smith, Tom, Mellor, Steve, Rare, Lawrence, Genton, Blaise
1998 The Effect of Distance from Home on Attendance at a Small Rural Health centre in Papua New Guinea. *International Journal of Epidemiology* 27:878-884.
- Nepal Census
2013 http://www.mohp.gov.np/english/files/new_publications/Nepal%20Population%20Report%202011.pdf, accessed November 11, 2014.
- O'Donnell, Owen
2007 Access to health care in developing countries: breaking down demand-side barriers. *Cad. Saúde Pública*, Rio de Janeiro, 23:12.
- Open Architecture Network
2012 OUTREACH: Mobile Health Clinics to Combat HIV/AIDS in Sub-Saharan Africa. <http://openarchitecturenetwork.org/projects/387>, accessed February 15, 2015.
- Pickering A. and Haddix McKay, K.H.
2007 Old and new barriers to family planning in Humla, Nepal: How the urban/rural divide has widened. *Contributions to Nepalese Studies*, 34(1): 229-245.
- Rebhan, David P.
2009 Health Care Utilization: Understanding and applying theories and models of health care seeking behavior. Case Western Reserve University. 1-19.
- Sanders, Catherine L. and Kimber H. McKay
2013 The Search For “Strong Medicine”: Pathways to Health care Development in Remote Nepal Using GIS. *Technology and Innovation* 15:109-124.
- Say, Lale, Raine, Rosalind
2007 A systematic review of the inequalities in the use of maternal health care in developing countries: examining the scale of the problem and the importance of context. *Bulletin of the World Health Organization*. 1-14.

Scheper-Hughes, Nancy

1992 *Death Without Weeping: The Violence of Everyday Life in Brazil*. Berkeley: University of California Press.

Scheper-Hughes, Nancy

1994 *Embodied Knowledge: Thinking with the Body in Critical Medical Anthropology*. In *Assessing Cultural Anthropology*, Rob Borofsky, ed. New York: McGraw-Hill, pp 229-242.

Schuler, SR, EN McIntosh, MC Goldstein, and BR Pande

1985 *Barriers to Effective Family Planning in Nepal*. *Studies in Family Planning* 16(5): 260-270.

Scott, James C.

2009 *The Art of Not Being Governed: An anarchist history of upland Southeast Asia*. New Haven: Yale University Press.

Singer, Merrill

1995 *Beyond the Ivory Tower: Critical Praxis in Medical Anthropology*. *Medical Anthropology Quarterly* 9(1):80-106.

Subedi, Madhusudan

2010 *Uterine Prolapse, Mobile Camp Approach and Body Politics in Nepal*. *Dhaulagiri Journal of Sociology and Anthropology* 4:21-40.

Sreeramareddy, Chandrashekhar T., Harsha Kumar, H.N., Sathian, Brijesh

2013 *Time Trends and Inequalities of Under-Five Mortality in Nepal: A Secondary Data Analysis of Four Demographic and Health Surveys between 1996 and 2011*. *PLOS ONE* 8(11):1-12.

Tipping, G., Segall M.

1995 *Health care seeking behaviour in developing countries: an annotated bibliography and literature review*. Sussex: Institute of Development Studies.

UCCC, Nepal

1991 *Self-Help Initiative Promotion Program, Thehe and Bargaun Village Development Committees, Humla District*. USCCN, Baluwatar, Kathmandu.

Unite for Sight

2000-2015(a) *The Complexities and Realities of Global Health*. New Haven, CT. <http://www.uniteforsight.org/global-health-university/complexities>. Accessed on January 13, 2015.

Unite for Sight

2000-2015(b): *Unique Barriers to Health Care for Women*. New Haven, CT <http://www.uniteforsight.org/women-children-course/barriers-women>. Accessed

on January 13, 2015.

United Nations Field Coordination Office (UNFCO)

2013 District Profile: Humla. Mid-Western Region, Nepalgunj, Nepal. 1-18.

Van Bodegom, May, Kuningas, Kaptijn, Thomese, Meij, Amankwa, Westendorp

2009 Chapter 4: Socio-economic status by rapid appraisal is highly correlated to mortality risks in rural Africa. Published in Transactions of the royal society of tropical medicine and hygiene 103:65-82.

Ward, Earlise C., Clark, Le Ondra, and Heidrich, Susan

2009 African American Women's Beliefs, Coping Behaviors, and Barriers to Seeking Mental Health Services. Qual Health Res. 19(11):1589-1601.

Ward, H., Mertens, T.E. and Thomas, C.

1997 Health seeking behavior and the control of sexually transmitted disease. Health Policy and Planning 12(1):19-28.

Wikipedia

2014 Humla District. http://en.wikipedia.org/wiki/Humla_District. Accessed on December 30, 2014.

Young, James Clay

1981 Medical Choice in a Mexican Village. New Brunswick, New Jersey. Rutgers University Press.