

11-2014

## A regional look at HANPP: human consumption is increasing, NPP is not

Steven W. Running

*University of Montana - Missoula*, [steven.running@umontana.edu](mailto:steven.running@umontana.edu)

Let us know how access to this document benefits you.

Follow this and additional works at: [https://scholarworks.umt.edu/decs\\_pubs](https://scholarworks.umt.edu/decs_pubs)

 Part of the [Earth Sciences Commons](#), and the [Environmental Sciences Commons](#)

---

### Recommended Citation

Steven W Running 2014 Environ. Res. Lett. 9 111003 doi:10.1088/1748-9326/9/11/111003

This Response or Comment is brought to you for free and open access by the Ecosystem and Conservation Sciences at ScholarWorks at University of Montana. It has been accepted for inclusion in Ecosystem and Conservation Sciences Faculty Publications by an authorized administrator of ScholarWorks at University of Montana. For more information, please contact [scholarworks@mail.lib.umt.edu](mailto:scholarworks@mail.lib.umt.edu).

A regional look at HANPP: human consumption is increasing, NPP is not

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 Environ. Res. Lett. 9 111003

(<http://iopscience.iop.org/1748-9326/9/11/111003>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 150.131.70.68

This content was downloaded on 11/06/2015 at 17:08

Please note that [terms and conditions apply](#).

## Perspective

# A regional look at HANPP: human consumption is increasing, NPP is not

**Steven W Running**

Numerical Terradynamic  
Simulation Group, University of  
Montana, Missoula Montana  
USA 59812

**Abstract**

Abdi *et al* (2014 *Environ. Res. Lett.* **9** 094003), have adapted the concept of comparing supply and demand of annual plant production known as human appropriation of net primary production (HANPP) to a region of the Sahel with rapid population growth. They found that HANPP more than doubled over the study period of 2000–2010, from 19% to 41%, suggesting increasing vulnerability of these populations to food insecurity.

Keywords: net primary production, food security, carbon cycle

Few issues are more fundamental to our continued presence on Earth than the sustainability of human consumption of biospheric resources. This question is often approached as an input–output or a supply–demand analysis of the terrestrial carbon cycle, with the input, or supply being NPP, and the output, or demand being human consumption of that NPP. At a global scale, HANPP or human appropriation of net primary production attempts to quantify what fraction of NPP is used or co-opted by humans, a broad definition that includes not only direct consumption for food, animal feed or fuel, but also NPP used for broader human benefit. The first estimates, by Vitousek *et al* (1986) and (1997), estimated that humanity was using about 32–40% of global NPP. More recent global estimates are 14–26% by Imhoff *et al* (2004), and 13–25% by Krausman *et al* (2013). These studies have the same policy motivation, to evaluate if humanity is approaching some sort of capacity, or limit in consumption of the biosphere.

The new paper by Abdi *et al* (2014) is a great example of analyzing HANPP at a very policy relevant regional scale. At issue here is comparing the biophysically controlled annual plant production (NPP) to the actual demand and consumption of that plant matter by the local population with the explicit goal of understanding if this represents a sustainable local carbon balance. The area of study encompassed 22 countries of sub-Saharan Africa, across a rather spectacular climatic gradient ranging from 100 to 600 mm of annual rainfall. These are developing countries with high reliance on their own land for the food, fibre and fuel (i.e. HANPP) for their societies. Abdi *et al* report that in this sub-Saharan study region 95% of domestic food consumption comes from crops grown in this area, there is very little import or export of crops, such that the region is quite self-sufficient, either by choice or more likely by economic necessity. Consequently, a rather complete carbon cycle supply–demand analysis is possible and politically important in this region. Spatially extensive satellite based measures of NPP were used to define the *supply* of NPP on each 1 km<sup>2</sup> of land. Abdi *et al* then computed NPP *demand* for human food, fuel, animal feed, forest burning for land clearing, and crop residues (see figure 1). The policy critical result was that while NPP supply was on aggregate stable during the study period from 2000 to 2010, population increased 31%, HANPP increased from 19% to 41% of NPP supply,



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.




Figure 1. The datastream used to compute a regional scale HANPP from Abdi *et al* 2014.

and per capita available NPP declined. Most significantly, on critical drought years, such as 2002, NPP supply dropped 9.7% below the decadal average, triggering food insecurity issues for 12 million people.

Other studies have analyzed capacities of the terrestrial carbon cycle for human use with different methodologies but similar goals. Terrestrial carbon accounting has typically been focused on the biophysical carbon source-sink dynamics of ecosystem NPP at regional scales (Houghton 2013). Ecosystem services for carbon are an attempt to blend biophysical analysis with socio-economic analysis (Costanza *et al* 2014). The optimum scale for these carbon cycle analyses varies, from a single product, to a local land-owner, to national policies and finally global carbon budgets. But the key concept is understanding the carbon cycle in an integrated, policy relevant way that evaluates the system capacity for NPP against a variety of demands and economic values. I have argued previously that at global scales NPP shows little capacity to be increased

(Running 2012). Globally, little additional land can be made available for food production beyond what already is being used (Lambin and Meyfroidt 2011). However, agricultural efficiency can be greatly improved in many regions, and in fact could successfully feed the projected global population of 9–10 billion in 2050 with proper policy choices (Mueller *et al* 2012). Mueller *et al* (2012) report that increasing NPP supply in sub-Saharan Africa should focus most by addressing agricultural nutrient deficiencies. If a changing climate brings more intense drought episodes, water management will be even more important. The largest unknown is new demands on terrestrial NPP, most notably large scale bioenergy, which may provide a direct competitor to food production in the regions that can least afford it (Smith *et al* 2012, Haberl *et al* 2013). Mueller *et al* (2012) and Abdi *et al* (2014) both recommend more efficient cooking stoves as a viable way to reduce bioenergy demand in this region.

The concept of HANPP originated as a global analysis, but regional applications such as the Abdi *et al* paper may in fact be more valuable for policy development by individual governments. The regional HANPP trend analysis can quantify increasing vulnerability of the local populations to any shortfall in NPP as demand approaches supply, an early warning system. Local policies to either increase NPP supply, or to reduce NPP demand may be better prioritized. Similar regional studies would be valuable in other parts of the world.

### Acknowledgements

The global NPP dataset used in this research resulted from multiple NASA Earth Science Division grants to SWR over the last 20 years.

### References

- Abdi A M, Seaquist J, Tenenbaum D E, Eklundh L and Ardo J 2014 The supply and demand of net primary production in the Sahel *Environ. Res. Lett.* **9** 094003
- Costanza R, deGroot R, Sutton P, van der Ploeg S, Anderson S, Kubiszewski I, Farber S and Turner R K 2014 Changes in the global value of ecosystem services *Glob. Environ. Change* **26** 152–8
- Imhoff M L, Bounoua L, Ricketts T, Loucks C, Harriss R and Laurance W T 2004 Global patterns in human consumption of net primary production *Nature* **429** 870–3
- Haberl H, Erb K, Krausmann F, Running S, Searchinger T D and Smith W K 2013 Bioenergy: how much can we expect for 2050? *Environ. Res. Lett.* **8** 031004
- Houghton R A 2013 Keeping management effects separate from environmental effects in terrestrial carbon accounting *Glob. Change Biol.* **19** 2609–12
- Krausmann F, Erb K-H, Gingrich S, Haberl H, Bondeau A, Gaube V, Lauk C, Plutzer C and Searchinger T D 2013 Global human appropriation of net primary production doubled in the 20th century *Proc. Natl. Acad. Sci. USA* **110** 10324–9
- Lambin E F and Meyfroidt P 2011 Global land use change, economic globalization, and the looming land scarcity *Proc. Natl. Acad. Sci. USA* **108** 3465–72
- Mueller N D, Gerber J S, Johnston M, Ray D K, Ramankutty N and Foley J A 2012 Closing yield gaps through nutrient and water management *Nature* **490** 254–7
- Running S W 2012 A measurable planetary boundary for the biosphere *Science* **337** 1458–9
- Smith W K, Zhao M and Running S W 2012 Global bioenergy capacity as constrained by observed biospheric productivity rates *BioScience* **62** 911–22
- Vitousek P M, Ehrlich P R, Ehrlich A H and Matson P A 1986 Human appropriation of the products of photosynthesis *BioScience* **36** 368–73
- Vitousek P M, Mooney H A, Lubchenco J and Melillo J M 1997 Human domination of Earth's ecosystems *Science* **277** 494–9